

Compendium of Measures to Prevent Disease Associated with Animals in Public Settings, 2023

The National Association of State Public Health Veterinarians Animal Contact Compendium Committee

Russell F. Daly, DVM, MPH, DACVPM (Co-chair); South Dakota State University, Brookings, SD

Abigail A. Mathewson DVM, MPH (Co-chair); New Hampshire Department of Health and Human Services, Concord, NH

Kerry R. Pride DVM, MPH, DACVPM; Montana Department of Public Health and Human Services, Helena, MT

Malia J. Ireland DVM, MPH, DACVPM; Minnesota Department of Health, St. Paul, MN

Sarah J. Bailey DVM, MPH, DACVPM; North Dakota Department of Agriculture, Bismarck, ND

Consultants to the Committee:

Karen Beck, North Carolina Department of Agriculture and Consumer Services, Raleigh, NC

Kathy Benedict, Centers for Disease Control and Prevention, Atlanta, GA

Marla Calico, International Association of Fairs and Expositions, Springfield, MO

Thomas Hairgrove, American Association of Extension Veterinarians, College Station, TX

Thomas Meehan, Association of Zoos and Aquariums, Silver Spring, MD

Christopher Olsen, American Veterinary Medical Association, Madison, WI

Stephan Schaeffbauer, US Department of Agriculture, South Saint Paul, MN

Sean Stapleton, Centers for Disease Control and Prevention, Atlanta, GA.

Address correspondence to Dr. Mathewson (abigail.a.mathewson@dhhs.nh.gov)

[I. Introduction](#)

[II. Methods](#)

[III. Background](#)

[A. Infectious diseases associated with animals in public settings](#)

[1. Diseases transmitted by direct or indirect animal contact](#)

[2. Diseases transmitted through droplets or aerosols](#)

[3. Factors influencing the risk of zoonotic disease transmission](#)

[B. Physical injuries caused by animals in public settings](#)

[IV. Recommendations for disease prevention](#)

[A. Overview](#)

[B. Applicable venues](#)

[C. Recommendations for local, tribal, state, and federal agencies](#)

[D. Recommendations for animal exhibitors and venue operators](#)

[1. Education](#)

[2. Facility design and use](#)

[3. Animal care and management](#)

[4. Birthing exhibits](#)

[5. Considerations regarding variant influenza](#)

[V. Summary – VI. Acknowledgements – VII. Footnotes](#)

[VIII. References](#)

[Appendix 1: Selected Zoonotic Diseases of Importance in Public Settings in the United States, 2023](#)

[Appendix 2: Animals in Public Settings: Recommendations for Venue Operators, Staff and Volunteers](#)

[Appendix 3: Handwashing Recommendations to Reduce Disease Transmission from Animals in Public Settings](#)

[Appendix 4: Guidelines for Exhibition of Animals in School and Childcare Settings](#)

[Appendix 5: Other Useful Disease Prevention Resources](#)

Preface

The Compendium of Measures to Prevent Disease Associated with Animals in Public Settings has been published by the NASPHV and the CDC since 2005.^[1-4] This compendium provides standardized recommendations for public health officials, veterinarians, animal venue operators, animal exhibitors, visitors to animal venues and exhibits, teachers, camp operators, and others concerned with control of disease and with minimizing health risks associated with animal contact in public settings. The report has undergone several revisions, and this document updates information provided in the 2017 compendium.^[4]

I. Introduction

Contact with animals in public settings (see **Table 1** for a comprehensive list) provides opportunities for entertainment and education. The National Association of State Public Health Veterinarians (NASPHV) understands the positive benefits of human-animal contact. However, an inadequate understanding among animal exhibitors and visitors regarding disease transmission and animal behavior can increase the likelihood of infectious disease exposures, injuries, and other health problems. Zoonotic diseases (i.e., zoonoses) are diseases shared between animals and people; many of these diseases are potentially transmitted from animals to people in public animal contact venues (**Appendix 1**). Of particular concern are instances in which zoonotic disease outbreaks result in numerous people becoming ill. For the 10-year period ending in 2020, 156 human infectious disease outbreaks involving animals in public settings resulted in 6,388 illnesses, 1,233 hospitalizations and 10 deaths.^[5] Such outbreaks, and additional infections that occur in these settings but are not related to outbreaks, have substantial medical, public health, legal, and economic effects.

Recommendations and best practices in this compendium were developed using a One Health approach with the goal of preventing zoonotic disease transmission between animals and people, and reducing zoonotic disease risks in environments where animals are displayed in public settings.⁶ Although completely eliminating risks from animal contact is not possible, this report provides recommendations for minimizing associated disease and injury. The NASPHV advises that local, tribal, territorial, and state public health, agricultural, animal health, wildlife, environmental, and educational agencies use these best practices to establish their own guidelines or regulations for reducing the risk for disease from human-animal contact in public settings. Public contact with animals is permitted in numerous types of venues (**Table 1**) and risk levels will vary by type of animals, type of encounters, pathogens, and other factors. Managers of these venues should use the information in this report in consultation with veterinarians, public health officials, state and local agriculture officials, or other professionals to evaluate and reduce risks for disease transmission for their specific venues.

Guidelines to reduce risks for disease from animals in health-care facilities, veterinary facilities, and various other occupational settings as well as from service and therapy animals have been developed.^[7-14] Although these settings are not specifically addressed here, the general principles and recommendations in this report are applicable. Recommendations to prevent illnesses associated with non-traditional pets (e.g., rodents, reptiles, fish, small mammals, backyard poultry) in private and public settings have recently been published.^[15] The NASPHV encourages managers of venues with these types of animals to reference that document as well.

II. Methods

The NASPHV periodically updates the recommendations to prevent disease associated with animals in public settings. To revise the 2017 compendium, the NASPHV Animal Contact Compendium Committee members and external consultants met virtually from September 12 through 16, 2022. The revision process included reviewing literature pertaining to outbreaks, diseases associated with animals in public settings, and prevention methods since the previous compendium was published. Specific input solicited from NASPHV members and committee consultants was also reviewed. This recent information was used to update the 2017 recommendations where applicable. A committee consensus was required to add or modify existing language or recommendations.

Table 1

Public Settings with Animals Present

- Agricultural Fairs
- Farms: educational, crop share, or otherwise open to public
- Agritourism venues
- Livestock birthing exhibits
- Petting zoos or animal displays
- Schools and childcare facilities
- Camps
- Zoological institutions
- Nature parks
- Pet stores
- Feed stores
- Live animal markets
- Animal swap meets
- Wildlife or animal photography settings
- Circuses and carnivals

III. Background

A. Infectious diseases associated with animals in public settings

1. Diseases transmitted by direct or indirect animal contact

One of the most common routes of disease transmission from animals to people is direct physical contact with the animal, which includes touching, holding, kissing, and being bitten, licked, and scratched. Disease transmission also occurs indirectly from an animal through contact with a surface contaminated by the animal's saliva, blood, urine, respiratory secretions, feces, or other bodily fluids or via contaminated food and beverages.

a. Enteric (intestinal) diseases

The considerable burden of enteric illness attributable to animal contact has been investigated in both the US and Canada.^[16,17] Pathogens with the highest proportion of cases attributable to animal contact were *Campylobacter* spp, nontyphoidal *Salmonella enterica* species, *Cryptosporidium* spp, non-O157 shiga-toxicogenic *Escherichia coli* (STEC), and STEC O157:H7. Annually in the US, close to half a million enteric illnesses are estimated to be due to animal contact, including 4,933 hospitalizations, and 76 deaths.^[16] In Canada, the annual estimate includes 6,000 illnesses, 488 hospitalizations and 12 deaths.^[17] More recent data estimates, for example, that 21% of all non-O157 STEC infections in the United States, 16% of *Campylobacter* infections, and 11-17% of *Salmonella* spp. infections are attributable to animal contact.^[18]

Enteric bacteria and parasites pose the most commonly encountered risk for human disease from animals in public settings.^[19] Numerous enteric disease outbreaks among visitors to fairs, farms, petting zoos, and other public settings are well documented.^[20-46] Cattle, sheep, or goats^[20, 22, 25, 26, 31-33, 35, 36, 39, 41, 43, 45, 46] have typically been identified as sources for infection. However, other domestic and wild animals are also established sources of enteric infections (**Table 2**) and outbreaks related to these animals occur.^[5] Animals that appear healthy can carry pathogens that cause illness in people. In all animal species that might harbor organisms that can cause enteric illness, it is possible for animals that appear healthy and clean to carry and shed the bacteria in their excreta, which can contaminate their fur, hair, feathers, scales, or skin. A small number of pathogens is often enough to cause illness.^[47-51]

Outbreaks and sporadic (i.e., those not linked to an outbreak) infections with *Salmonella* spp in particular have been associated with animal contact from a wide variety of species. Animals that have been implicated as sources of outbreaks of salmonellosis include, but are not limited to: poultry,^[21, 52-63] reptiles,^[38, 64-78] amphibians^[79-82] and other non-traditional pets such as hedgehogs,^[83, 84] hamsters, mice, guinea pigs^[85-89] and aquatic species such as African dwarf frogs.^[79, 81] Contact with animals causing illness has occurred at feed stores, schools, daycare facilities, fairs, agritourism venues, petting zoos, nursing homes, carnivals, pet or retail stores, aquatic exhibits and public farm visits. A compendium of measures to prevent zoonotic diseases associated with non-traditional pets was recently published by NASPHV.^[15] This publication details the multiple pathogens, animal sources, exposures settings, and risk behaviors involved in these infections and provides detailed recommendations for the prevention of illness from non-traditional pets in private and public settings.

Outbreaks of enteric infections are not the only source of animal contact-related illness. Sporadic cases also occur in public and private settings.^[77, 91-99] A 2019 study of non-outbreak illnesses associated with animal agriculture found that the agricultural exposure for 10% of cases was a fair, petting zoo, agritourism farm or other public animal agriculture venue.^[100] In fact, public animal agricultural settings caused more sporadic illnesses than outbreak cases during the 5-year study. For STEC cases, 28% had visited a public animal contact setting, including six cases who developed hemolytic uremic syndrome.

(1) Animals shedding enteric pathogens.

Animals carrying enteric pathogens frequently have no signs of illness but can still shed the organisms in

Table 2

Animals recognized as sources for enteric disease outbreaks in public settings

- Cattle
- Goats
- Sheep
- Live Poultry (chickens, ducks, geese, turkeys)^[21, 52-59]
- Rodents^[85-89]
- Hedgehogs^[83, 84]
- Reptiles^[38, 64-78]
- Amphibians^[79-82]
- Wild animals^[90]

feces.^[101] Removing ill animals, especially those with diarrhea, from public contact is necessary, but this step alone is not sufficient to protect the health of people and other animals. The fact that some pathogens can be shed intermittently and survive for months or years in the environment,^[102-106] as well as the limitations of laboratory testing, makes attempts to identify and remove infected animals unreliable as means of eliminating the risk for transmission. Antimicrobial treatment cannot reliably eliminate infection or prevent shedding, and it does not protect against reinfection. Antimicrobial use in animals can also prolong shedding and contribute to antimicrobial resistance.^[107-109] Disease transmission at animal exhibits can be influenced by multiple factors. Stress induced by transportation, confinement, physical crowding, and increased handling increases the likelihood of animals shedding pathogens.^[110-116] Commingling increases the probability that the shedding pathogens will infect other animals.^[117] Young animals, which are frequently included in public settings and educational programs for children, have a higher prevalence of shedding enteric pathogens such as STEC than do mature animals.^[118-120]

(2) Direct transmission of enteric pathogens from animals to people.

Enteric pathogens are primarily transmitted by the fecal-oral route. Because animal fur, hair, feathers, scales, skin, and saliva harbor fecal organisms,^[121] transmission can occur when people pet, touch, kiss, hold, feed, or are licked by animals and bacteria are ingested. In addition, illness has resulted from fecal contamination of food^[29, 122] and beverages^[24, 123-131] at animal contact settings.^[128-131]

(3) Indirect transmission and environmental exposures to enteric pathogens.

Disease transmission can occur in the absence of direct animal contact if a pathogen is present in the environment, which occurs when animals or animal excreta, fur, hair, feathers, scales, or skin are present. Exposure to contaminated materials such as animal bedding, environmental surfaces, clothing, and shoes has been associated with transmission of pathogens.^[34, 38, 40, 81, 132, 133] Outbreaks of enteric illness have been associated with exposure to environments after animals were removed,^[134] dust in the environment,^[133] touching or stepping in manure or shavings,^[37] contact with manure on a fence,^[27] haybales,^[46] manure piles,^[135] and falling down or sitting on the ground in a petting zoo.^[37] In an outbreak of STEC in 2004, the outbreak strain was isolated from shavings collected from a baby stroller and from the shoes of petting zoo visitors.^[37] Pet food and treats, which may be present in public settings such as pet stores, fairs, and school classrooms, have been confirmed as sources of human salmonellosis in several instances.^[136-142] House flies may also contaminate surfaces with pathogens.^[143] Furthermore, enteric pathogens can persist in contaminated environments for long periods. For example, STEC can survive in soil for months.^{[27, 40, 101, 102, 104, 106, 133], a}

(4) Populations with additional risks

While anyone can get sick due to enteric or other pathogens from animals in public settings, some groups are more likely to become ill or to have severe consequences of infection (**Table 3**)^[144]. The recommendations that follow are particularly important for these groups, including providing education about the risks.

b. Internal parasites

Animal parasites can infect people who ingest materials contaminated with animal feces or who ingest or otherwise come into contact with contaminated soil. Exposure to parasites can occur in public settings as well as other places where animals are present. Pollution of public settings and agritourism venues with feral and domestic cat feces has led to outbreaks of toxoplasmosis.^[147-149] Other parasites of concern include *Toxocara* (roundworms) and *Ancylostoma* (hookworms) which can lead to larval migrans.^[147, 148, 150-153] The risk of *Baylisascaris* (raccoon roundworm) exposure from infected animals in public settings is low; however, environmental contamination with raccoon feces is possible.^[154] While human infection with *Giardia* spp. is also considered an environmental hazard from wildlife, advances in genetic testing suggests that

Table 3

Populations with additional risks

- **Young children** (i.e., < 5 years of age) are at greater risk for acquiring enteric pathogens due to their behaviors. Closer physical contact with animals and hand-to-face contact were more common in children than in adults during petting zoo visits.^[144, 145]
- **People with weakened immune systems** such as young children and adults 65 and older, have an increased risk for developing severe illness, compared with healthy individuals outside these groups, when they do become infected.^[146]
- Despite their more frequent exposure to livestock, **exhibitors and/or employees** have also become infected with enteric pathogens in outbreaks at fairs.^[40]

domestic animals in public settings do not pose a significant risk of transmitting pathogenic giardiasis to people.^[155-158]

c. Animal bites and scratches

(1) Rabies

People who have contact with rabid mammals can be exposed to rabies virus via a bite or when mucous membranes or open wounds come into contact with infected saliva or nervous tissue. In the United States, multiple rabies exposures have occurred requiring extensive public health investigations and medical follow-up; no deaths have occurred. While the number of exposures by type of setting is not known, it is estimated that thousands of people receive rabies postexposure prophylaxis after being exposed to rabid or potentially rabid animals or animal carcasses. Species involved in reported exposures have included bats, raccoons, cats, goats, bears, sheep, horses, foxes, and dogs, at venues including an urban public park,^[159] a pet store,^[160] a county fair,^[161, 162] petting zoos,^[163, 164] schools,^[162] rodeo events,^[162] a horse show,^[165] and summer camps.^[166] Public health challenges associated with possible mass rabies exposures include difficulty in identifying and contacting individuals who are potentially at risk, correctly assessing exposure risks, and providing timely medical prophylaxis when indicated. Human infection with rabies virus is almost always fatal once clinical signs of rabies appear in the infected person. Prompt assessment and appropriate treatment after exposure are critical.^[167]

(2) Other bite-related and scratch-related infections.

Infections from animal bites and scratches are common; some might require extensive treatment or hospitalization. Bacterial pathogens associated with animal bites include *Pasteurella* spp, *Francisella tularensis*,^[168, 169] *Staphylococcus* spp, *Streptococcus* spp, *Capnocytophaga canimorsus*, *Bartonella* spp (etiologic agents of cat scratch disease), and *Streptobacillus moniliformis* (the etiologic agent of rat bite fever).^[170] Some monkey species (especially macaques) can be infected with B virus. Infected monkeys might have no clinical signs or have mild oral lesions. Fatal meningoencephalitis has been reported in human patients infected via monkey bites or by exposure to bodily fluids.^[171, 172]

d. Skin infections

Contact with animals in public settings can also result in human skin infection. Cases of ringworm have been reported among animal exhibitors.^[173] Infection with the parapox virus that causes contagious ecthyma or sore mouth in sheep and goats (and orf in humans) has developed in children after contact with sheep in a public setting.^[174] Transmission of other pox viruses to people in public settings also has been described, including cowpox virus in a circus animal keeper,^[175] cowpox virus in people who handled pet rats at a pet store,^[176] and monkeypox among people who contacted infected prairie dogs at a childcare center.^[177, 178] Contact with aquatic animals and their environment has also been implicated in cutaneous infections,^[179] such as *Mycobacterium marinum* infections in people who owned or had cleaned fish tanks.^[180, 181]

e. External parasites

Fleas, ticks, and mites can be acquired by people who interact with animals in public settings.^[182] *Sarcoptes scabiei* is a skin mite with different host-specific variants that infest people and animals, including swine, dogs, cats, foxes, and cattle.^[183, 184] Although human infestation with animal mites is typically self-limiting, skin irritation and itching might occur for multiple days and can be difficult to diagnose.^[184, 185]

2. Diseases transmitted through droplets or aerosols

Generation of infectious droplets or aerosols and subsequent contamination of the environment is an important risk for indirect transmission of disease in public settings. These droplets or aerosols can include infectious agents from animals' respiratory tracts, reproductive fluids, or other sources. Cleaning procedures (e.g., pressure washing^[12, 186]) or dust raised in animal environments, including dust generated from activities such as sweeping and leaf blowing, can lead to infectious aerosols in the immediate environment and surrounding areas.

a. Influenza

Transmission of influenza A viruses between people and animals has increasingly important implications for human-animal interactions in public settings. Influenza viruses that normally circulate in pigs are called variant influenza viruses when they are found in people.^[187] Although pigs with influenza can become ill, it has also been shown that apparently healthy pigs can carry influenza viruses.^[188] Sporadic cases and small clusters of human infections with variant influenza viruses have been reported since the 1970s.^[188-195] The CDC continues to provide

information about these cases and assists in response to their identification when necessary (https://gis.cdc.gov/grasp/fluview/Novel_Influenza.html).

Cases of human infection with variant influenza viruses from pigs have been associated with direct or indirect exposure to swine at agricultural fairs^[196-198] and many of these infections occurred in children who reported direct contact with swine at agricultural fairs. To manage the risk of variant influenza at these events, a specialized guidance document has been developed, “Measures to Minimize Influenza Transmission at Swine Exhibitions,” and should be followed.^[195]

Although viruses that normally circulate in birds (avian influenza A viruses) usually do not infect humans, rare cases of human infection with these viruses have been reported.^[194] Transmission of human influenza viruses from people to swine^[200, 201] and other species also has been reported. For example, in 1998, a new strain of influenza A (H3N2) virus derived from human, avian, and classical swine influenza A viruses emerged and became established in swine.^[202] Swine and poultry species should be kept separated because of the potential for transmission of avian influenza viruses to pigs and, conversely, swine influenza to birds, especially turkeys. Adaptation of avian viruses to pigs and genetic mixing of viruses in pigs can create novel viruses with the potential to infect and spread in the human population.^[203]

b. Tuberculosis

Tuberculosis, although rare, can be a concern in certain animal settings; however, the risk is primarily for close contact between handlers and certain animal species,^[204-206] including elephants.^[204, 207, 208] Guidelines have been developed regarding diagnosis, treatment and management of tuberculosis in elephants, as well as for the removal of tuberculosis-infected animals from public settings.^[209]

c. Q fever

Live-birthing exhibits, usually involving cattle, pigs, goats, or sheep, are popular at agricultural fairs and farm visits. Although members of the public do not typically have direct contact with animals during birthing, some infections, such as Q fever, can be acquired by inhalation. Cases of illness related to Q fever have been linked to viewing of animal births.^{[210], b}

The causative agent of Q fever is the *Coxiella burnetii* bacterium; goats, sheep, and cattle are the most frequently implicated animal sources of human infections in the United States.^[211] Although *C. burnetii* infection can cause abortion in animals, it is often subclinical. High numbers of organisms shed in reproductive tissues, and fluids can become aerosolized during birthing, and inhalation of aerosolized organisms leads to infection in people. Most individuals exposed to *C. burnetii* develop an asymptomatic infection, but clinically apparent illness can range from an acute influenza-like illness to life-threatening endocarditis, as well as premature birth, stillbirth, and miscarriage in pregnant women.^[212] In 1999, an outbreak of Q fever involving 95 confirmed cases of the disease and 41 hospitalizations was linked to goats and sheep giving birth at petting zoos in indoor shopping malls in Canada.^b Another Q fever outbreak, in which > 30 human cases were reported in the Netherlands, was associated with public lamb-viewing days at a sheep farm in 2009.^[210]

Leptospirosis, listeriosis, brucellosis, and chlamydiosis are other serious zoonotic diseases that can be acquired through contact with aborted fetuses, newborn animals, reproductive tissues, or associated fluids associated with animal birthing.^[90]

d. *Chlamydia psittaci* infections

Chlamydia psittaci infections are usually acquired from psittacine birds (e.g. parrots) and cause respiratory disease in people. Guidelines are available for control of this infection.^[213] Cases of human psittacosis have occurred among staff members at a zoological garden,^[214] among people exposed to an aviary in a church,^[215] and among pet store staff and visitors.^[43] On rare occasions, chlamydial infections acquired from sheep and birds have resulted in human maternal and fetal illness and death.^[213, 216-219]

3. Factors influencing the risk of zoonotic disease transmission

a. Handwashing

Handwashing following contact with animals has been associated with decreased rates of illness during disease outbreaks associated with animals in public settings. Centers for Disease Control and Prevention (CDC) was prompted to establish recommendations for enteric disease prevention associated with farm animal contact after two outbreaks of STEC infections in 2000 in Pennsylvania and Washington.^[220] Risk factors identified in the Pennsylvania outbreak were contact with cattle and inadequate handwashing. It was found that handwashing facilities were limited and not configured for children.^[41]

In 1996, an outbreak of salmonellosis at a Colorado zoo resulted in 65 cases of the disease (primarily among children) associated with touching a wooden barrier around a temporary Komodo dragon exhibit. Children who were not ill were significantly more likely to have washed their hands after visiting the exhibit than children who were ill.^[38]

Handwashing has proven to be a mitigating factor in illness in outbreaks of STEC and *Salmonella* infections associated with touching animals, barriers, and sawdust.^[30, 32, 38] c The likelihood of illness was higher for people who reported feeding animals and not washing hands prior to eating and drinking when compared with those who did. Creating a lather decreased the likelihood of illness for individuals who used soap and water for handwashing; however, drying hands on clothing increased the likelihood of illness.

Interventions that have been shown to improve hand hygiene compliance include having venue staff provide verbal reminders about hand hygiene to guests before they leave the animal area, use of larger signs with more prominent messages combined with staff actively offering hand sanitizer to visitors,^[221] and having a staff member present within or at the exit to the animal contact area.^[222] Although the use of hand sanitizers (with an alcohol concentration of 60% to 95%) can be effective at killing pathogens, it should be noted that washing hands with soap and water is still preferred because hand sanitizers do not work equally well for all classes of pathogens and do not work well when hands are heavily soiled or greasy.^[223]

b. Facility design

The layout and maintenance of facilities and animal exhibits can increase or decrease the risk for infections.^[224] Factors that increase this risk include inadequate handwashing facilities,^[162] inappropriate flow of visitors, and incomplete separation between animal exhibits and food preparation and consumption areas.^[34, 43, 225] Other factors that increase risk include structural deficiencies associated with temporary food service facilities, contaminated or inadequately maintained drinking water systems, and poorly managed sewage or manure containment and disposal processes.^[38, 129-131, 133, 226] In one of the largest waterborne disease outbreaks in the United States (1999),^[129, 130] approximately 800 suspected cases of infection with STEC, *Campylobacter* spp, or both were identified among attendees at a New York county fair. In that outbreak, unchlorinated water supplied by a shallow well was used by food vendors to make beverages and ice.^[130]

Temporary and seasonal animal exhibits and activities are particularly vulnerable to design flaws.^[30, 38] Animal displays or petting zoos added to attract visitors to zoos, festivals, roadside attractions, farm stands, farms where people can pick their own produce, feed stores, and Christmas tree lots are examples. In 2004 and 2005, separate outbreaks of STEC occurred at seasonal state fairs in North Carolina and Florida. Both of these outbreaks involved exposure to vendor-run temporary petting zoos.^[30] Inadequate handwashing facilities were reported for a temporary exhibit in British Columbia, Canada, where childcare facility and school field trips to a pumpkin patch with a petting zoo resulted in STEC infections.^[43] Running water and signs recommending handwashing were not available, and alcohol-containing hand sanitizers were placed at a height that was unreachable for some children.

Venues not designed for or accustomed to public events, such as working farms, wildlife rehabilitation facilities, animal adoption events, and animal shelters, might be less likely to have facilities adequately designed to accommodate visitors and to reduce the risk of exposure to zoonotic disease agents. Limitations that might lead to increased infection risk include lack of or inadequate handwashing stations and dedicated food service areas and inappropriate traffic flow patterns. Public access to animal waste areas in these venues might also be problematic.^[134]

c. Food contamination

Contamination of food products, beverages, or food preparation areas by people after animal contact has previously resulted in outbreaks.^[24, 29, 123, 129, 162] Contamination from inadequate sanitation (e.g., of hands, utensils, or equipment) can occur during food preparation or consumption. Venues in which such food contamination contributed to human illness include summer camps^[29] and an apple orchard.^d Large, multistate foodborne outbreaks of salmonellosis have been attributed to food preparers having had contact with live poultry prior to handling food products and subsequently contaminating those products.^[21, 227] Additionally, consumption of food in an animal environment has been associated with illnesses. In a 2015 outbreak of STEC infections at a dairy event in Washington, crude attack rates were higher for individuals who were involved in activities where food was served in an animal barn.^[134] Purchase of food at a farm visit^[220] and the consumption of sticky foods^[122] (e.g., ice cream and cotton candy) have also been associated with STEC-related illnesses.

d. Other factors influencing disease transmission

Events at which people have prolonged close contact with animals, such as day camps and livestock exhibitions, pose a unique challenge with regard to disease prevention. Examples of events where prolonged contact has led to

illness include an outbreak of STEC infections that occurred at a day camp where prolonged contact with livestock was encouraged.^[228]

Failure to properly implement disease-prevention recommendations has also contributed to recurrent outbreaks. Following an outbreak of cryptosporidiosis with 31 ill students at an educational farm program in Minnesota, specific recommendations (including use of coveralls and rubber boots when handling calves, supervised handwashing, and provision of hand sanitizer) were provided to teachers but were inadequately implemented.^[36] A subsequent outbreak occurred several months later, with 37 additional illnesses.^[36] Handwashing facilities and procedures were still inadequate, and coveralls and boots that were used were found to be dirty, cleaned infrequently, and handled without subsequent handwashing.

Other disease outbreaks have resulted from contaminated animal products used during school activities. Salmonellosis outbreaks associated with dissection of owl pellets in classes have occurred^[229]; in one such outbreak, risk factors for infection included inadequate handwashing, use of food service areas for the activity, and improper cleaning of contact surfaces. Students in a middle school science class were among those infected in a multi-state salmonellosis outbreak associated with frozen rodents sold as snake food.^[87]

B. Physical injuries caused by animals in public settings

Although infectious diseases are the most commonly reported health problems associated with animals in public settings, injuries caused by animals are also commonly reported, and these can result in infection as well as trauma. For example, dog bites are an important community problem for which specific guidelines have been written.^[230] Injuries associated with animals in public settings include bites, kicks, falls, scratches, stings, crushing of extremities, and being pinned between an animal and a fixed object. Serious and fatal injuries have been associated with various venues and species including commercial stables (interaction with horses),^[231] animal sanctuaries (tigers),^[232] petting zoos (llamas),^[233] photo opportunities (tigers and bison),^[232, 234] schools (snakes),^[235] animal safaris (camels),^[236] and dog parks (dogs).^[237]

IV. Recommendations for Disease Prevention

A. Overview

Information, publications, and reports from multiple organizations were used to create the recommendations in this document using a One Health approach.^[238-240] No US federal laws address the risk for transmission of pathogens at venues where animals and the public come into contact. However, some states regulate certain disease prevention practices such as the provision of handwashing stations and animal vaccination requirements.^[241, 242]

Certain federal agencies and associations in the United States have developed standards, recommendations, and guidelines for reducing health risks associated with animal contact by the public. The Association of Zoos and Aquariums has accreditation standards requiring training of staff on the risks of zoonotic diseases, including those associated with public contact.^[243] The USDA licenses and inspects certain animal exhibits in accordance with the Animal Welfare Act^[244]; although these inspections primarily address humane treatment of animals, they also impact animal health and public safety. In 2001, CDC first issued recommendations to reduce the risk of infection with enteric pathogens associated with farm visits, and has also provided recommendations for preventing transmission of *Salmonella* spp from reptiles, amphibians, and live poultry to people.^[15, 62, 71, 72, 81, 220, 245] The Association for Professionals in Infection Control and Epidemiology and the Animal-Assisted Interventions Working Group have developed guidelines to address risks associated with the use of animals in health-care settings.^[10, 13] The NASPHV has developed guidance and compendia of measures to reduce risks for human exposure to *C. psittaci*, rabies virus, *C. burnetii*, novel influenza A viruses, zoonotic pathogens arising from non-traditional pets such as live poultry, reptiles, and rodents, and zoonotic pathogens that veterinary personnel might be exposed to in an occupational setting.^[12, 15, 213, 246-248]

Studies^[145, 144, 221, 222, 249] in multiple localities have suggested that the recommendations provided in the present compendium are not completely implemented by members of the public and managers or employees of animal contact venues. Stakeholders should strive to achieve comprehensive implementation of the recommendations in this compendium, to help ensure that visitors can stay healthy and reduce the risk of zoonotic disease transmission while enjoying animals.

B. Applicable venues

The recommendations in this report were developed for settings in which contact with animals or their environments is possible. These settings include farm visits, agritourism venues, petting zoos, school field trips, camps, agricultural fairs, feed stores, wildlife sanctuaries, animal swap meets, childcare centers and schools, and other settings (**Table 1**). Contact with animals in public settings should only occur where measures are in place to reduce the potential for disease transmission or injuries. Incidents or problems should be investigated, documented, and reported.

C. Recommendations for local, tribal, state, territorial, and federal agencies

Agencies should encourage or require oversight to ensure compliance with recommendations at animal contact

venues. The recommendations should be tailored to specific settings using a One Health approach and incorporated into best practices, protocols, and regulations developed at the state or local, or tribal level. Additional research should be conducted regarding the risk factors and effective prevention and control methods for health issues associated with animal contact. Additionally, communication and cooperation to ensure public health and safety extends beyond human, animal, and environmental health agencies and should include additional stakeholders such as professional associations, schools, private companies, and industry groups.

1. Dissemination of recommendations

This compendium should be disseminated to cooperative extension personnel, venue operators, farms that host public events, veterinarians, schools and daycares, associations and industry groups, public health officials, and others associated with managing animals in public settings. Development of a complete list of public animal contact venues within a jurisdiction is encouraged to facilitate dissemination of these recommendations. Agencies should disseminate educational and training materials to venue operators and other stakeholders. Sample materials are available in a variety of media in the NASPHV Animals in Public Settings Toolkit, which is available electronically.^[250]

2. Investigating and reporting outbreaks

To evaluate and improve these recommendations, surveillance activities for human infections associated with animal contact should be enhanced. Agencies should take the following steps:

- Conduct thorough epidemiological investigations of outbreaks using a One Health approach across human, animal, and environmental health sectors.
- Follow appropriate protocols for collection and laboratory testing of samples from people, animals, and the environment, including molecular subtyping of pathogen isolates.
- Include questions on disease report forms and outbreak investigation questionnaires about exposure to animals and their husbandry, environments, products, feed, and veterinary care.
- Report outbreaks to state public health departments.
- Local, tribal, territorial, and state public health departments should also report all outbreaks of enteric infections resulting from animal contact to the CDC through the National Outbreak Reporting System (www.cdc.gov/nors/).

D. Recommendations for animal exhibitors and venue operators

Staff and visitor education, attention to hygiene, and appropriate facility design as well as proper care and monitoring of animals and their enclosures are essential components for reduction of risks associated with animal contact in public settings. It is important to be aware of and follow local, tribal, state, and federal regulations regarding animals in public settings. Animal exhibitors and venue operators and staff should implement and encourage record keeping practices that support disease outbreak investigations, should they be needed to identify where animals originated and when animals have been moved between multiple public settings.^[251]

1. Education

Awareness of zoonotic disease risk is protective against illness in outbreaks.^[37] Therefore, educating visitors to public animal contact venues about the risk for transmission of diseases from animals to humans is a potential disease-prevention measure. Education is important not only at traditional animal venues like petting zoos, but also at farms and other venues where live animals are sold or distributed to the public. Even in well-designed venues with operators who are aware of the risks for disease, outbreaks and injuries can occur when visitors do not understand the risks and therefore are less likely to apply disease-prevention measures. This is especially important for animals considered to have a high risk of transmitting disease to humans (e.g., reptiles and live poultry); hatcheries, agricultural feed stores, and pet stores are other venues where particular efforts should be made to provide information to potential owners.^[15] Evidence-based prevention messages and free educational materials are available in multiple formats and in multiple languages on the CDC Healthy Pets, Healthy People website (www.cdc.gov/healthypets).

a. Operators and staff

Operators and staff should be aware that certain human populations are more likely than others to develop serious illness from pathogens transmitted in animal contact settings (see **Table 3**). Individuals considered to be at high risk for serious illness, in addition to thorough and frequent handwashing, should take heightened precautions or avoid animal exhibits and their environments.

Venue operators and staff (all individuals involved with animal contact activity in any public setting) should take the following steps for public health and safety:

- Become familiar with and implement the recommendations in this compendium.
- Consult with veterinarians, state and local agencies, and cooperative extension personnel on implementation of the recommendations.
- Become knowledgeable about the risks for disease and injury associated with animals and be able to explain risk-reduction measures to staff members and visitors.
- Be aware of human populations at high risk for disease and injury (see **Table 3**).
- Understand which animals pose a high risk for causing disease and injury within the venue.

Each of the following aspects should be taken into consideration in facility design and operation, educational messaging, and animal care and management:

- Direct public contact with ill animals is inappropriate for any audience.
- Children < 5 years of age (**Table 3**) should not have direct contact with animals that are considered likely to carry zoonotic pathogens (e.g., preweaned calves, reptiles, live poultry, or other species with demonstrated risks).

- Children < 5 years of age are also at high risk for disease and injury from contact with other animals and should be supervised at all times to discourage hand-to-mouth activities (e.g., nail biting and thumb sucking), contact with manure, and contact with soiled bedding.

- Individuals > 65 years of age and those with weakened immune systems (e.g., people with HIV-AIDS, without a functioning spleen, or receiving immunosuppressive treatment) also have a high risk of developing serious illness from contact with animals carrying zoonotic diseases (**Table 3**).

- Pregnant people are at risk of stillbirth, miscarriage, and preterm delivery from certain pathogens that might be present in animal contact settings.

- Direct contact with venomous or otherwise dangerous animals (e.g., venomous reptiles, nonhuman primates, or certain carnivores and other rabies reservoir species) should be completely prohibited (see the Animal Care and Management section for more information on these species)

- Live animals should not be given as prizes at fairs, carnivals, or other events.

- Ensure that visitors receive educational instructions before entering an exhibit, including information that animals can cause injuries or carry pathogens that can cause serious illness, along with recommended prevention measures (**Figure 1**; **Appendix 2**).^[1-4]

- Provide information in a simple and easy-to-understand format that is age appropriate and language appropriate.
- Provide information in multiple formats (e.g., signs [including explanatory images and diagrams if appropriate], stickers, handouts, and verbal




Figure 1—Suggested sign or handout for use in safety education of visitors entering animal areas of petting zoos or other exhibits.^[250]



Figure 2—Suggested sign to encourage compliance with handwashing procedures as a means of reducing the possible spread of infectious disease.^[252]

information) and languages.

- Provide information to people arranging school field trips or classroom exhibits so they can educate participants and parents before the visit.
- Encourage compliance by the public with risk reduction recommendations, especially compliance with handwashing procedures as visitors exit animal areas (**Figure 2; Appendix 3**).^[1-4, 253]
- Ensure compliance with licensing and registration requirements under the Animal Welfare Act per USDA guidelines for dealers, exhibitors, transporters, and researchers.^[244]
- Comply with local, tribal, territorial, and state requirements for reporting animal bites or other injuries.

b. Visitors

Visitors to animal exhibits and those participating in interaction activities of any kind should be presented with effective educational messages aimed at ensuring compliance with the following recommendations:

- Be aware that the risks of, and severity of disease associated with, animal contact are higher among people of certain age groups and health conditions (**Table 3**).
- Supervise children properly at all times while in the presence of animals and areas with animal waste; prevent inappropriate contact with animals and sitting or playing on the ground.
- Refrain from eating, drinking, or other hand-to-mouth activities in the presence of animals.
- Practice proper hand hygiene, including washing hands immediately upon exit of the animal area and before any hand-to-mouth activity or eating is done.^[252]
- Practice proper hand hygiene after any contact with shoes, strollers, or clothing that might have come in contact with animals, their waste, or their bedding.^[252]
- Report any animal bites or injuries promptly to the venue operator and to authorities per local, tribal, or state law.
- Understand that certain diseases shared between animals and people can also pass from people to animals.^[254]

2. Facility design and use


Venues should be divided into 3 types of areas: animal areas (where animal contact is possible or encouraged), transition areas (located at entrances and exits to animal areas), and nonanimal areas (where animals are not permitted, with the exception of service animals; **Figure 3**).

a. Layout and traffic patterns

(1) Animal area considerations

The design of facilities and animal pens should minimize the risk associated with animal contact (**Figure 3**), including limiting direct contact with manure and encouraging handwashing (**Appendix 3**). The design of facilities or contact settings might include double barriers to prevent contact with animals or contaminated surfaces except in specified animal interaction areas. Contact with fecal material or soiled bedding in animal pens increases risk of exposure to pathogens, and facility designs and policies should limit or prevent this type of exposure, especially to individuals who might be at high risk for infection.

Investigations of previous outbreaks have revealed that temporary exhibits are often not designed appropriately. Common problems include inadequate barriers, floors and other surfaces that are difficult to keep clean and disinfect, insufficient plumbing, lack of signs regarding potential health risks and risk prevention measures, and inadequate handwashing facilities.^[30, 37, 38, 122] Specific recommendations might be necessary for certain other settings, such as schools and childcare facilities (**Appendix 4**).^[1-4]



Recommendations for animal areas are as follows:

- Do not allow consumption of food or beverages in animal areas.
- Do not allow toys, pacifiers, spill-proof cups, baby bottles, strollers or similar items to enter animal areas.
- Individuals utilizing wheelchairs or other mobility devices should consider how these items can be adequately cleaned and disinfected following their visit.
- Prohibit smoking and other tobacco product use in animal areas.
- Do not allow children to sit or play on the ground in animal areas or on manure piles. If hands become soiled, supervise handwashing immediately.
- For areas where animal contact is encouraged, a 1-way flow of visitors is recommended, with separate entrance and exit points (**Figure 3**).
- Control visitor traffic to prevent overcrowding.
- Ensure that animal feed bowls or bins and water are not accessible to the public.
- Allow the public to feed animals only in circumstances where contact with animals is controlled (e.g., with barriers).
- Do not provide animal feed in containers that can be eaten by people (e.g., ice cream cones) to decrease the possibility of children eating food that has come into contact with animals.
- Promptly remove manure and soiled animal bedding from exhibit areas.
- Assign trained staff members to encourage appropriate human-animal interactions, to identify and reduce potential risks for patrons, and to process reports of injuries and exposures.
- Ensure that visitors do not have access to animals that are not part of the defined interaction area, especially in on-farm visit situations.
- Store animal waste and specific tools for waste removal (e.g., shovels and pitchforks) in designated areas that are restricted from public access.
- Avoid transporting manure and soiled bedding through nonanimal areas or transition areas. If this is unavoidable, take precautions to prevent spillage, or remove manure when the public is not present.
- Where feasible, clean and disinfect the animal area (e.g., flooring and railings) as necessary.
- Provide adequate ventilation for animals^[243] and people, but avoid creating air movement that distributes dust, which may contain contaminants.
- Minimize the use of animal areas for public activities (e.g., weddings and dances). If areas previously used for animals must be used for public events, they should be cleaned and disinfected, particularly if food or beverages are served.
- For bird encounter exhibits, refer to the NASPHV's psittacosis compendium^[213] for recommendations regarding disease prevention and control.
- Visitors to aquatic touch tank exhibits should be advised not to participate if they have open wounds. Handwashing stations and signs should be provided as for other venues.
- When using animals or animal products (e.g., pelts, fecal material, or owl pellets) for educational purposes, use them only in designated areas, and not in school cafeterias or other areas where food and beverages are stored, prepared, served, or consumed.
- When animals are in school or childcare settings, specific areas must be designated for animal contact (**Appendix 4**). These areas must be thoroughly cleaned after use. Parents and guardians should be informed of the presence of animals as well as the benefits and potential risks associated with animals in schools and childcare settings.
- Immersion exhibits (where members of the public enter into the animal space) present additional opportunities for transmission of infectious agents. Entry into these spaces can lead to increased contamination of clothes, shoes, and other items, therefore increasing risk for disease. Lack of barriers between animals and people also increases the risk for injury. These exhibits heighten the need for supervision and awareness by venue operators and attendees.

(2) Transition area considerations.

The following steps are recommended for management of transition areas between non-animal and animal areas. Establishing transition areas through which visitors pass when entering and exiting animal areas is critical. The transition areas should be designated as clearly as possible, even if they are conceptual rather than physical (**Figure 3**).

Entrance transition areas should be designed to facilitate education:

- Position venue staff members (when available), post signs, and otherwise notify visitors that they are entering an animal area and that there are risks associated with animal contact, including who might be at highest risk for illness. **(Figure 1)**.
- Instruct visitors not to eat, drink, smoke, place their hands in their mouth, or use bottles or pacifiers while in the animal area.
- Establish storage or holding areas for strollers and related items (e.g., wagons and diaper bags).

Exit transition areas should be designed to facilitate handwashing **(Appendix 3)**:

- Post signs or otherwise instruct visitors to wash their hands when leaving the animal area **(Figure 2)**.
- Provide accessible handwashing stations for all visitors, including children and people with disabilities **(Figure 3)**.
- Position venue staff members near exits to encourage compliance with proper handwashing.
- Post signs or otherwise instruct visitors to exercise proper handwashing when handling shoes, clothing, strollers, wheelchairs, and other mobility devices that might have come in contact with animal bedding or waste.

(3) Nonanimal area considerations.

Recommendations for nonanimal areas are as follows:

- Do not permit animals, except for service animals, in nonanimal areas.
- Restrict storage, preparation, serving, and consumption of food and beverages to nonanimal areas.
- Provide handwashing facilities and display handwashing signs where food or beverages are served **(Figure 2; Appendix 3)**.
- Separation of food from animal contact areas is of particular importance to farm visit venues; this includes food tasting, distribution of food samples, and consumption of beverages, snacks, or meals.

b. Cleaning and disinfection

Cleaning and disinfection practices should be tailored to the specific situation. For example, most parasitic pathogens, such as *Cryptosporidium parvum*, are resistant to most disinfectants. When a particular organism has been identified, additional guidance regarding specific disinfectants can be found in other resources.^[255] General recommendations are that all surfaces should be cleaned thoroughly to remove organic matter before disinfection. Prompt, safe removal of fecal matter reduces the risk of infection. Disinfectants, such as bleach and quaternary ammonium, should be used in accordance with the manufacturer label. Most compounds require > 10 minutes of contact time with a contaminated surface to achieve the desired result. Animals should be removed during the cleaning process and should not reenter the area until after disinfected surfaces have been thoroughly dried.

Venue operators should strive to develop an integrated pest management program to eliminate or reduce the risk of exposure to pathogens carried by pests. Carriers of concern include flies, mosquitos, ticks, and fleas as well as rodents and wild birds.

c. Unpasteurized food and products

Unpasteurized or raw dairy products (e.g., milk, cheese, and yogurt) and unpasteurized cider or juices are potential sources of foodborne pathogens.^[123, 126, 146, 256] The sale or distribution of such products should be prohibited.

d. Drinking water

Local public health authorities should inspect drinking water systems before use. Only potable water should be used for consumption by animals and people. Backflow prevention devices should be installed between outlets in livestock areas and water lines supplying other areas on the grounds. The use of outdoor hoses should be minimized, and hoses should not be left on the ground. Hoses that are accessible to the public should be labeled to indicate the water is not for human consumption. Operators and managers of facilities in settings where treated municipal water is not available should ensure that a safe water supply (e.g., bottled water) is available.

3. Animal care and management

a. Selection of animals for use in public settings

The risk for disease or injury from animal contact can be reduced by carefully managing animal use. The following recommendations should be considered for management of animals in contact with the public:

- Direct contact with some animals is inappropriate in public settings, depending on expected audiences. Use of preweaned calves, reptiles, amphibians, and live poultry (including chicks) is not appropriate in nursing homes, schools, daycares, or other venues where groups at high risk for serious infection are expected to be present; contact with other young ruminants such as lambs or goat kids is also of increased concern in such settings.
- Animals showing signs of illness are not appropriate for use in public settings.
- Direct contact with species known to serve as reservoirs for rabies virus (e.g., bats, raccoons, skunks, foxes, and coyotes) should not be permitted.
- Certain nonhuman primates are of particular concern because of the types of pathogens they can transmit to people, such as B virus.^[257]
- Because of their strength, unpredictability, or ability to produce venom, certain domestic, exotic, or wild animals should be prohibited from exhibition settings where a reasonable possibility of animal contact exists. Species of primary concern include certain nonhuman primates, certain carnivores (e.g., lions, tigers, ocelots, wolves and wolf hybrids, and bears), and venomous species (e.g., some reptiles and invertebrates).

b. Routine animal care

Venue operators and staff should monitor animals daily for signs of illness and ensure that animals receive appropriate veterinary medical care. Ill or injured animals, animals known to be infected with a zoonotic pathogen, and animals from herds with a recent history of abortion, diarrhea, or respiratory disease should not be exhibited. To decrease shedding of pathogens, animals should be housed in a manner to minimize stress, injury, and overcrowding.

c. Veterinary medical care and animal health

Venue operators should retain and use the services of a licensed veterinarian. Regular inspection while animals are present in the venue is a critical component of monitoring health. When necessary, Certificates of Veterinary Inspection from an accredited veterinarian should be up to date according to local or state requirements for animals in public settings. Preventive care, including vaccination appropriate for the species, should be provided with appropriate input from the attending veterinarian. Parasite control methods should be used for animals in public settings to reduce the risk of human exposure to flea and tick-borne diseases.

(1) Vaccination against rabies virus.

All animals should be housed in a manner that reduces potential exposure to wild animals that may serve as rabies virus reservoirs. Mammals should also be up to date for rabies vaccinations according to current recommendations.^[246] These steps are particularly critical in areas where rabies is endemic and in venues where human-animal contact is encouraged or possible. Because of the extended incubation period for rabies, unvaccinated mammals should be vaccinated ≥ 1 month before they have contact with the public. If no licensed rabies vaccine exists for a particular species that is used in a setting where public contact occurs, consultation with a veterinarian regarding extralabel use of rabies vaccine is recommended.^[246] Mammals that are too young to be vaccinated should be used in exhibit settings only if additional restrictive measures are available to reduce risks (e.g., using only animals that were born to vaccinated mothers and housed in a manner to avoid rabies exposure). In animal contact settings, rabies testing should be considered for animals that die suddenly.

(2) Vaccination against zoonotic pathogens.

While vaccines against certain enteric pathogens (e.g., *Salmonella* spp and STEC) are available for specific animal species, insufficient evidence currently exists to support the use of these products to reduce transmission of disease to people in public settings.^[258] More research is necessary and encouraged before firm recommendations can be made. Vaccination of slaughter-class animals before displaying them at fairs might not be feasible because of the slaughter withdrawal period that is needed when certain vaccines are used. Vaccination of animals in public settings for diseases other than rabies should be done in consultation with a licensed veterinarian and tailored to the animal species and nature of public interaction.

(3) Testing for zoonotic pathogens.

Routine screening for zoonotic diseases is not recommended, except for *C. psittaci* infection in bird encounter exhibits^[213] and tuberculosis in elephants^[204] and primates.^[259] Screening tests are available for some enteric pathogens; however, the interpretation of test results can be problematic. Shedding can be intermittent, and negative results do not indicate an animal was not shedding an organism at an earlier time or will not start shedding in the near future. There is no established guidance for management of animals with positive test

results, and inappropriate interpretation might lead to unnecessary treatments, quarantine, or euthanasia.

4. Birthing exhibits

Animal birthing exhibits are increasingly popular. However, it is important for organizers and attendees to understand that animals such as goats, sheep, and cattle giving birth might be shedding pathogens, such as *C. burnetii*, *Brucella* spp, *Leptospira* spp, and *L. monocytogenes*.^[90] Organizers should be aware of the following steps to reduce the risk of disease transmission:

- Ensure that the public has no contact with newly born animals or birthing byproducts (e.g., the placenta).
- Ensure that attendees and staff who are particularly vulnerable to zoonotic diseases (e.g., pregnant women, people with cardiac valvular disease and other heart conditions, and people with weakened immune systems) and the parents of small children understand the risks of attending or working at these exhibits.
- Thoroughly clean and disinfect the birthing area after each birth, and use appropriate personal protective equipment (PPE), safety precautions, and disposal methods for discarding waste products.
- If abortions or stillbirths occur, the exhibit should be closed; operators should work with their veterinarians to determine the cause of abortions or stillbirths.
- Birthing events should be held outdoors or in well-ventilated areas to reduce the risk for human exposure to aerosolized pathogens. Additional information is available electronically in the CDC fact sheet on Q fever safety at livestock birthing exhibits.^[260]

5. Considerations regarding variant influenza

In response to influenza A variant virus outbreaks associated with swine at agricultural fairs since 2011, the following prevention strategies have been recommended^[195]:

- All people should take routine preventive actions (e.g., practice appropriate hand hygiene) at fairs to reduce potential influenza virus transmission between pigs and people.
- People at increased risk of serious influenza-related complications should avoid exposure to pigs at fairs.
- Shortening the time that animals are on exhibit has been shown to dramatically reduce the risk of influenza transmission in swine, lowering the risk of human illness due to zoonotic influenza.^[261]
- Measures should be taken to reduce the presence of pigs with clinical signs of disease at these events.

Potential strategies to mitigate the risk for intraspecies and interspecies transmission of influenza viruses at agricultural fairs include shortening the swine exhibition period,^[262] consulting with a veterinarian to determine whether vaccination of swine against influenza is appropriate, keeping swine and poultry separate,^[263] cleaning and disinfecting sorting boards, and providing for ≥ 7 days between exhibitions to reduce the risk of spreading influenza.^[195] More detailed and current recommendations for fairs have been published and can also be found at the NASPHV website.^[195]

V. Summary

Contact and interaction with animals in public settings can be a valuable means of education and entertainment. People who provide these opportunities to the public as well as those attending such venues should be aware of the potential health risks associated with such venues and understand that even apparently healthy animals can transmit pathogens. The recommendations included in this compendium will help venue operators, staff, and attendees reduce the risk for injury and zoonotic disease transmission in these settings.

VI. Acknowledgments

The authors thank Rochelle Medford, Colin Basler, Ann Carpenter, Kate Varela, Natalie Wendling, and Casey Barton Behravesh from CDC's One Health Office for their work in revising and updating references and materials in this compendium.

VII. Footnotes

a. Durso LM, Keen JE, Bauer N, et al. Assessment of three re- mediation strategies for reduction of Shiga-toxigenic *Escherichia coli* (STEC) O157 in naturally contaminated soil (abstr), in *Proceedings. Annu Meet Inst Food Technol* 2007. Available at: <https://www.ars.usda.gov/research/publications/publication/?seqNo115=207142>. Accessed Mar 6, 2023.

b. Milford F, Vibien A, Lambert L, et al. Large Q-fever outbreak related to exposure to petting zoos in two shopping malls (oral presentation). *51st Annu Conf Dis Nat Transmissible Man*, Austin, Tex, May–June 2001.

c. Chertow D, Gupta S, Ginzl D, et al. Outbreak of *Escherichia coli* O157:H7 related to direct and indirect animal contact in petting zoos—Florida, 2005 (abstr), in *Proceedings. 55th Annu Epidemic Intell Serv Conf* 2006;36.

d. Saupe A, Fowler H, Anderston F, et al. *E. coli* O111 and *Cryptosporidium* infections associated with raw apple cider at an apple orchard (poster presentation). *OutbreakNet Meet*, Atlanta, Ga, August 2012.

VIII. References

1. National Association of State Public Health Veterinarians, *Compendium of measures to prevent disease associated with animals in public settings, 2013*. *J Am Vet Med Assoc*, 2013. 243(9): p. 1270-88.
2. National Association of State Public Health Veterinarians, *Compendium of measures to prevent disease associated with animals in public settings, 2005*. *MMWR Recomm Rep*, 2005. 54(RR-4): p. 1-12.
3. National Association of State Public Health Veterinarians, Centers for Disease Control and Prevention, *Compendium of measures to prevent disease associated with animals in public settings, 2011*: *MMWR Recomm Rep*, 2011. 60(RR-04): p. 1-24.
4. National Association of State Public Health Veterinarians, *Compendium of Measures to Prevent Disease Associated with Animals in Public Settings, 2017*. *J Am Vet Med Assoc*, 2017. 251(11): p. 1268-1292.
5. Centers for Disease Control and Prevention. *National Outbreak Reporting System (NORS)*. Atlanta, Georgia: U.S. Department of Health and Human Services, CDC. Download date: 18 Nov 2022.
6. Centers for Disease Control and Prevention. *One Health*. Available at: www.cdc.gov/onehealth/. Accessed May 16, 2023.
7. Hardin, P., J. Brown, and M.E. Wright, *Prevention of transmitted infections in a pet therapy program: An exemplar*. *Am J Infect Control*, 2016. 44(7): p. 846-50.
8. Duncan, S.L., *APIC State-of-the-Art Report: the implications of service animals in health care settings*. *Am J Infect Control*, 2000. 28(2): p. 170-80.
9. Guay, D.R., *Pet-assisted therapy in the nursing home setting: potential for zoonosis*. *Am J Infect Control*, 2001. 29(3): p. 178-86.
10. Lefebvre, S.L., et al., *A veterinary perspective on the recently published guidelines for animal-assisted interventions in health-care facilities*. *J Am Vet Med Assoc*, 2008. 233(3): p. 394-402.
11. Murthy, R., et al., *Animals in healthcare facilities: recommendations to minimize potential risks*. *Infect Control Hosp Epidemiol*, 2015. 36(5): p. 495-516.
12. Williams, C.J., et al., *Compendium of Veterinary Standard Precautions for Zoonotic Disease Prevention in Veterinary Personnel: National Association of State Public Health Veterinarians: Veterinary Infection Control Committee 2015*. *J Am Vet Med Assoc*, 2015. 247(11): p. 1252-77.
13. Sehulster, L., et al., *Guidelines for environmental infection control in health-care facilities. Recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee (HICPAC)*. *MMWR Recomm Rep*, 2003. 52(RR-10): p. 1-42.
14. National Institute for Occupational Safety and Health. *Veterinary Safety and Health*. Available at: www.cdc.gov/niosh/topics/veterinary/default.html. Accessed Mar 6, 2023.
15. Varela, K., et al., *A Review of Zoonotic Disease Threats to Pet Owners: A Compendium of Measures to Prevent Zoonotic Diseases Associated with Non-Traditional Pets: Rodents and Other Small Mammals, Reptiles, Amphibians, Backyard Poultry, and Other Selected Animals*. *Vector Borne Zoonotic Dis*, 2022. 22(6): p. 303-360.
16. Hale, C.R., et al., *Estimates of enteric illness attributable to contact with animals and their environments in the United States*. *Clin Infect Dis*, 2012. 54 Suppl 5: p. S472-9.
17. Murray, R., et al., *Estimates of the burden of illness for eight enteric pathogens associated with animal contact in Canada*. *Epidemiol Infect*, 2017. 145(16): p. 3413-3423.
18. Beshearse, E., et al., *Attribution of Illnesses Transmitted by Food and Water to Comprehensive Transmission Pathways Using Structured Expert Judgment, United States*. *Emerg Infect Dis*, 2021. 27(1): p. 182-195.
19. LeJeune, J.T. and M.A. Davis, *Outbreaks of zoonotic enteric disease associated with animal exhibits*. *J Am Vet Med Assoc*, 2004. 224(9): p. 1440-5.
20. Moller-Stray, J., et al., *Two outbreaks of diarrhoea in nurseries in Norway after farm visits, April to May 2009*. *Euro Surveill*, 2012. 17(47).
21. Loharikar, A., et al., *Four multistate outbreaks of human Salmonella infections associated with live poultry contact, United States, 2009*. *Zoonoses Public Health*, 2012. 59(5): p. 347-54.
22. Utsi, L., et al., *Cryptosporidiosis outbreak in visitors of a UK industry-compliant petting farm caused by a rare Cryptosporidium parvum subtype: a case-control study*. *Epidemiol Infect*, 2016. 144(5): p. 1000-9.
23. Adams, N.L., et al., *Shiga Toxin-Producing Escherichia coli O157, England and Wales, 1983-2012*. *Emerg Infect Dis*, 2016. 22(4): p. 590-7.

24. Brooks, J.T., et al., *An outbreak of Salmonella serotype Typhimurium infections with an unusually long incubation period*. Foodborne Pathog Dis, 2012. 9(3): p. 245-8.
25. Ihekweazu, C., et al., *Large outbreak of verocytotoxin-producing Escherichia coli O157 infection in visitors to a petting farm in South East England, 2009*. Epidemiol Infect, 2012. 140(8): p. 1400-13.
26. McGuigan, C.C., K. Steven, and K.G. Pollock, *Cryptosporidiosis associated with wildlife center, Scotland*. Emerg Infect Dis, 2010. 16(5): p. 895-6.
27. Lanier, W.A., et al., *Outbreak of Shiga-toxigenic Escherichia coli O157:H7 infections associated with rodeo attendance, Utah and Idaho, 2009*. Foodborne Pathog Dis, 2011. 8(10): p. 1131-3.
28. Centers for Disease, C. and Prevention, *Notes from the field: Escherichia coli O157:H7 gastroenteritis associated with a State Fair - North Carolina, 2011*. MMWR Morb Mortal Wkly Rep, 2012. 60(51-52): p. 1745-6.
29. Centers for Disease, C.P., *Cryptosporidiosis outbreak at a summer camp—North Carolina, 2009*. MMWR Morb Mortal Wkly Rep 2011. 60(918– 922).
30. Centers for Disease, C. and Prevention, *Outbreaks of Escherichia coli O157:H7 associated with petting zoos--North Carolina, Florida, and Arizona, 2004 and 2005*. MMWR Morb Mortal Wkly Rep, 2005. 54(50): p. 1277-80.
31. Warshawsky, B., et al., *Outbreak of Escherichia coli O157:H7 related to animal contact at a petting zoo*. The Canadian journal of infectious diseases = Journal canadien des maladies infectieuses, 2002. 13(3): p. 175-181.
32. Smith, K.E., et al., *Outbreaks of enteric infections caused by multiple pathogens associated with calves at a farm day camp*. Pediatr Infect Dis J, 2004. 23(12): p. 1098-104.
33. Shukla, R., et al., *Escherichia coli O157 infection associated with a farm visitor centre*. Commun Dis Rep CDR Rev, 1995. 5(6): p. R86-90.
34. Sayers, G.M., et al., *Cryptosporidiosis in children who visited an open farm*. Commun Dis Rep CDR Rev, 1996. 6(10): p. R140-4.
35. Pritchard, G.C., et al., *Verocytotoxin-producing Escherichia coli O157 on a farm open to the public: outbreak investigation and longitudinal bacteriological study*. Veterinary Record, 2000. 147(10): p. 259-264.
36. Kiang, K.M., et al., *Recurrent outbreaks of cryptosporidiosis associated with calves among students at an educational farm programme, Minnesota, 2003*. Epidemiol Infect, 2006. 134(4): p. 878-86.
37. Goode, B., et al., *Outbreak of escherichia coli O157: H7 infections after Petting Zoo visits, North Carolina State Fair, October-November 2004*. Arch Pediatr Adolesc Med, 2009. 163(1): p. 42-8.
38. Friedman, C.R., et al., *An outbreak of salmonellosis among children attending a reptile exhibit at a zoo*. J Pediatr, 1998. 132(5): p. 802-7.
39. Evans, M.R. and D. Gardner, *Cryptosporidiosis outbreak associated with an educational farm holiday*. Commun Dis Rep CDR Rev, 1996. 6(3): p. R50-1.
40. Durso, L.M., et al., *Shiga-toxigenic Escherichia coli O157:H7 infections among livestock exhibitors and visitors at a Texas County Fair*. Vector Borne Zoonotic Dis, 2005. 5(2): p. 193-201.
41. Crump, J.A., et al., *An Outbreak of Escherichia coli O157:H7 Infections among Visitors to a Dairy Farm*. New England Journal of Medicine, 2002. 347(8): p. 555-560.
42. Chapman, P.A., J. Cornell, and C. Green, *Infection with verocytotoxin-producing Escherichia coli O157 during a visit to an inner city open farm*. Epidemiol Infect, 2000. 125(3): p. 531-6.
43. David, S.T., et al., *Petting zoo-associated Escherichia coli O157:h7--secondary transmission, asymptomatic infection, and prolonged shedding in the classroom*. Can Commun Dis Rep, 2004. 30(20): p. 173-80.
44. Halsby, K.D., et al., *Healthy animals, healthy people: zoonosis risk from animal contact in pet shops, a systematic review of the literature*. PLoS One, 2014. 9(2): p. e89309.
45. Lange, H., et al., *Second outbreak of infection with a rare Cryptosporidium parvum genotype in schoolchildren associated with contact with lambs/goat kids at a holiday farm in Norway*. Epidemiol Infect, 2014. 142(10): p. 2105-13.
46. Nichols, M.C., et al., *Agritourism and Kidding Season: A Large Outbreak of Human Shiga Toxin-Producing Escherichia coli O157 (STEC O157) Infections Linked to a Goat Dairy Farm-Connecticut, 2016*. Front Vet Sci, 2021. 8: p. 744055.
47. Robinson, D.A., *Infective dose of Campylobacter jejuni in milk*. Br Med J (Clin Res Ed), 1981. 282(6276): p. 1584.
48. Chappell, C.L., et al., *Cryptosporidium parvum: intensity of infection and oocyst excretion patterns in healthy volunteers*. J Infect Dis, 1996. 173(1): p. 232-6.
49. Tilden, J., Jr., et al., *A new route of transmission for Escherichia coli: infection from dry fermented salami*. Am J Public Health, 1996. 86(8): p. 1142-5.
50. Tuttle, J., et al., *Lessons from a large outbreak of Escherichia coli O157:H7 infections: insights into the infectious dose and method of widespread contamination of hamburger patties*. Epidemiol Infect, 1999. 122(2): p. 185-92.
51. Mody RK, O.R.C., Griffin PM, *E. coli diarrheal diseases*, in Heymann DL, ed. *Control of communicable diseases manual*. 2014, American Public Health Association: Washington, DC. p. 158–172.
52. Centers for Disease, C. and Prevention, *Three outbreaks of salmonellosis associated with baby poultry from three hatcheries--United States, 2006*. MMWR Morb Mortal Wkly Rep, 2007. 56(12): p. 273-6.
53. Gaffga, N.H., et al., *Outbreak of salmonellosis linked to live poultry from a mail-order hatchery*. N Engl J Med, 2012.

- 366(22): p. 2065-73.
54. Loharikar, A., et al., *Outbreak of human Salmonella Typhimurium infections linked to contact with baby poultry from a single agricultural feed store chain and mail-order hatchery, 2009*. *Pediatr Infect Dis J*, 2013. 32(1): p. 8-12.
 55. Basler, C., et al., *Notes from the field: multistate outbreak of human salmonella infections linked to live poultry from a mail-order hatchery in Ohio--February-October 2014*. *MMWR Morb Mortal Wkly Rep*, 2015. 64(9): p. 258.
 56. Nakao, J.H., et al., *'One Health' investigation: outbreak of human Salmonella Braenderup infections traced to a mail-order hatchery - United States, 2012-2013*. *Epidemiol Infect*, 2015. 143(10): p. 2178-86.
 57. Basler, C., et al., *Multistate outbreak of human Salmonella infections linked to live poultry from a mail-order hatchery in Ohio--March-September 2013*. *MMWR Morb Mortal Wkly Rep*, 2014. 63(10): p. 222.
 58. Centers for Disease Control and Prevention, *Notes from the field: Multistate outbreak of Salmonella infantis, newport, and lille infections linked to live poultry from a single mail-order hatchery in Ohio--March-September, 2012*. *MMWR Morb Mortal Wkly Rep*, 2013. 62(11): p. 213.
 59. Centers for Disease Control and Prevention, *Notes from the field: multistate outbreak of Salmonella Altona and Johannesburg infections linked to chicks and ducklings from a mail-order hatchery - United States, February-October 2011*. *MMWR Morb Mortal Wkly Rep*, 2012. 61(11): p. 195.
 60. Centers for Disease Control and Prevention, *Multistate outbreaks of Salmonella infections associated with live poultry--United States, 2007*. *MMWR Morb Mortal Wkly Rep*, 2009. 58(2): p. 25-9.
 61. Centers for Disease Control and Prevention, *Salmonella hadar associated with pet ducklings--Connecticut, Maryland, and Pennsylvania, 1991*. *MMWR Morb Mortal Wkly Rep*, 1992. 41(11): p. 185-7.
 62. Centers for Disease Control and Prevention, *Salmonellosis associated with chicks and ducklings--Michigan and Missouri, Spring 1999*. *MMWR Morb Mortal Wkly Rep*, 2000. 49(14): p. 297-9.
 63. Centers for Disease Control and Prevention, *Salmonella serotype Montevideo infections associated with chicks--Idaho, Washington, and Oregon, spring 1995 and 1996*. *MMWR Morb Mortal Wkly Rep*, 1997. 46(11): p. 237-9.
 64. Harris, J.R., et al., *Multistate outbreak of Salmonella infections associated with small turtle exposure, 2007-2008*. *Pediatrics*, 2009. 124(5): p. 1388-94.
 65. Centers for Disease Control and Prevention, *Notes from the field: outbreak of salmonellosis associated with pet turtle exposures--United States, 2011*. *MMWR Morb Mortal Wkly Rep*, 2012. 61(4): p. 79.
 66. Bosch, S., R.V. Tauxe, and C.B. Behravesh, *Turtle-Associated Salmonellosis, United States, 2006-2014*. *Emerg Infect Dis*, 2016. 22(7): p. 1149-55.
 67. Gambino-Shirley, K., et al., *Notes from the Field: Four Multistate Outbreaks of Human Salmonella Infections Linked to Small Turtle Exposure - United States, 2015*. *MMWR Morb Mortal Wkly Rep*, 2016. 65(25): p. 655-6.
 68. Walters, M.S., et al., *Outbreaks of Salmonellosis From Small Turtles*. *Pediatrics*, 2016. 137(1).
 69. Basler, C.A., et al., *Notes From the Field: Multistate Outbreak of Human Salmonella Poona Infections Associated With Pet Turtle Exposure—United States, 2014*. *American Journal of Transplantation*, 2016. 16.
 70. Whitten, T., et al., *Reptile-associated salmonellosis in Minnesota, 1996-2011*. *Zoonoses Public Health*, 2015. 62(3): p. 199-208.
 71. Centers for Disease Control and Prevention, *Multistate outbreak of human Salmonella infections associated with exposure to turtles--United States, 2007-2008*. *MMWR Morb Mortal Wkly Rep*, 2008. 57(3): p. 69-72.
 72. Centers for Disease Control and Prevention, *Reptile-associated salmonellosis--selected states, 1998-2002*. *MMWR Morb Mortal Wkly Rep*, 2003. 52(49): p. 1206-9.
 73. Centers for Disease Control and Prevention, *Turtle-associated salmonellosis in humans--United States, 2006-2007*. *MMWR Morb Mortal Wkly Rep*, 2007. 56(26): p. 649-52.
 74. Centers for Disease Control and Prevention, *Multistate outbreak of human Salmonella typhimurium infections associated with pet turtle exposure - United States, 2008*. *MMWR Morb Mortal Wkly Rep*, 2010. 59(7): p. 191-6.
 75. Harris, J.R., et al., *Recent multistate outbreaks of human salmonella infections acquired from turtles: a continuing public health challenge*. *Clin Infect Dis*, 2010. 50(4): p. 554-9.
 76. Lamm, S.H., et al., *Turtle-associated salmonellosis. I. An estimation of the magnitude of the problem in the United States, 1970-1971*. *Am J Epidemiol*, 1972. 95(6): p. 511-7.
 77. Mermin, J., et al., *Reptiles, amphibians, and human Salmonella infection: a population-based, case-control study*. *Clin Infect Dis*, 2004. 38 Suppl 3: p. S253-61.
 78. Altman, R., et al., *Turtle-associated salmonellosis. II. The relationship of pet turtles to salmonellosis in children in New Jersey*. *Am J Epidemiol*, 1972. 95(6): p. 518-20.
 79. Mettee Zarecki, S.L., et al., *US outbreak of human Salmonella infections associated with aquatic frogs, 2008-2011*. *Pediatrics*, 2013. 131(4): p. 724-31.
 80. Bartlett, K.H., T.J. Trust, and H. Lior, *Small pet aquarium frogs as a source of Salmonella*. *Appl Environ Microbiol*, 1977. 33(5): p. 1026-9.
 81. Centers for Disease Control and Prevention, *Multistate outbreak of human Salmonella typhimurium infections associated with aquatic frogs - United States, 2009*. *MMWR Morb Mortal Wkly Rep*, 2010. 58(51): p. 1433-6.

82. Centers for Disease Control and Prevention, *Notes from the field: update on human Salmonella Typhimurium infections associated with aquatic frogs--United States, 2009-2011*. MMWR Morb Mortal Wkly Rep, 2011. 60(19): p. 628.
83. Anderson, T.C., et al., *Multistate Outbreak of Human Salmonella Typhimurium Infections Linked to Pet Hedgehogs - United States, 2011-2013*. Zoonoses Public Health, 2017. 64(4): p. 290-298.
84. Centers for Disease Control and Prevention, *Notes from the field: Multistate outbreak of human Salmonella typhimurium infections linked to contact with pet hedgehogs - United States, 2011-2013*. MMWR Morb Mortal Wkly Rep, 2013. 62(4): p. 73.
85. Centers for Disease Control and Prevention, *Notes from the field: Infections with Salmonella I 4,[5],12:i:- linked to exposure to feeder rodents - United States, August 2011-February 2012*. MMWR Morb Mortal Wkly Rep, 2012. 61: p. 277.
86. Bartholomew, M.L., et al., *Multistate outbreak of Salmonella enterica serotype enteritidis infection associated with pet guinea pigs*. Vector Borne Zoonotic Dis, 2014. 14(6): p. 414-21.
87. Fuller, C.C., et al., *A multi-state Salmonella Typhimurium outbreak associated with frozen vacuum-packed rodents used to feed snakes*. Zoonoses Public Health, 2008. 55(8-10): p. 481-7.
88. Harker, K.S., et al., *An outbreak of Salmonella Typhimurium DT191a associated with reptile feeder mice*. Epidemiol Infect, 2011. 139(8): p. 1254-61.
89. Swanson, S.J., et al., *Multidrug-resistant Salmonella enterica serotype Typhimurium associated with pet rodents*. N Engl J Med, 2007. 356(1): p. 21-8.
90. Stuenkel S. Small Ruminants and Zoonotic Infections: Live or Dead—Direct or Indirect. In: Sing A, editor. Zoonoses - Infections Affecting Humans and Animals: Focus on Public Health Aspects. Dordrecht: Springer Netherlands; 2015. p. 69-101.
91. Voetsch, A.C., et al., *Risk factors for sporadic Shiga toxin-producing Escherichia coli O157 infections in FoodNet sites, 1999-2000*. Epidemiol Infect, 2007. 135(6): p. 993-1000.
92. Younus, M., et al., *The role of exposures to animals and other risk factors in sporadic, non-typhoidal Salmonella infections in Michigan children*. Zoonoses Public Health, 2010. 57(7-8): p. e170-6.
93. Kassenborg, H.D., et al., *Farm visits and undercooked hamburgers as major risk factors for sporadic Escherichia coli O157:H7 infection: data from a case-control study in 5 FoodNet sites*. Clin Infect Dis, 2004. 38 Suppl 3: p. S271-8.
94. Cummings, K.J., et al., *Farm animal contact as risk factor for transmission of bovine-associated Salmonella subtypes*. Emerg Infect Dis, 2012. 18(12): p. 1929-36.
95. O'Brien, S.J., G.K. Adak, and C. Gilham, *Contact with farming environment as a major risk factor for Shiga toxin (Vero cytotoxin)-producing Escherichia coli O157 infection in humans*. Emerg Infect Dis, 2001. 7(6): p. 1049-51.
96. Haack, J.P., et al., *Escherichia coli O157 exposure in Wyoming and Seattle: serologic evidence of rural risk*. Emerg Infect Dis, 2003. 9(10): p. 1226-31.
97. Hunter, P.R., et al., *Sporadic cryptosporidiosis case-control study with genotyping*. Emerg Infect Dis, 2004. 10(7): p. 1241-9.
98. Roy, S.L., et al., *Risk factors for sporadic cryptosporidiosis among immunocompetent persons in the United States from 1999 to 2001*. J Clin Microbiol, 2004. 42(7): p. 2944-51.
99. Friedman, C.R., et al., *Risk factors for sporadic Campylobacter infection in the United States: A case-control study in FoodNet sites*. Clin Infect Dis, 2004. 38 Suppl 3: p. S285-96.
100. Klumb, C.A., J.M. Scheftel, and K.E. Smith, *Animal agriculture exposures among Minnesota residents with zoonotic enteric infections, 2012-2016*. Epidemiol Infect, 2020. 148: p. e55.
101. Keen, J.E., et al., *Shiga-toxigenic Escherichia coli O157 in agricultural fair livestock, United States*. Emerg Infect Dis, 2006. 12(5): p. 780-6.
102. Kudva, I.T., K. Blanch, and C.J. Hovde, *Analysis of Escherichia coli O157:H7 survival in ovine or bovine manure and manure slurry*. Appl Environ Microbiol, 1998. 64(9): p. 3166-74.
103. LeJeune, J.T., T.E. Besser, and D.D. Hancock, *Cattle water troughs as reservoirs of Escherichia coli O157*. Appl Environ Microbiol, 2001. 67(7): p. 3053-7.
104. Maule, A., *Survival of verocytotoxigenic Escherichia coli O157 in soil, water and on surfaces*. Symp Ser Soc Appl Microbiol, 2000(29): p. 71S-78S.
105. Rahn, K., et al., *Persistence of Escherichia coli O157:H7 in dairy cattle and the dairy farm environment*. Epidemiol Infect, 1997. 119(2): p. 251-9.
106. Randall, L.P., C. Wray, and R.H. Davies, *Survival of verocytotoxin-producing Escherichia coli O157 under simulated farm conditions*. Vet Rec, 1999. 145(17): p. 500-1.
107. Berge, A.C., D.A. Moore, and W.M. Sischo, *Field trial evaluating the influence of prophylactic and therapeutic antimicrobial administration on antimicrobial resistance of fecal Escherichia coli in dairy calves*. Appl Environ Microbiol, 2006. 72(6): p. 3872-8.
108. Beraud, R., et al., *Comparison of the selection of antimicrobial resistance in fecal Escherichia coli during enrofloxacin administration with a local drug delivery system or with intramuscular injections in a swine model*. Can J Vet Res, 2008.

- 72(4): p. 311-9.
109. Johns, I., et al., *Antimicrobial resistance in faecal Escherichia coli isolates from horses treated with antimicrobials: a longitudinal study in hospitalised and non-hospitalised horses*. *Vet Microbiol*, 2012. 159(3-4): p. 381-9.
 110. Corrier, D.E., C.W. Purdy, and J.R. DeLoach, *Effects of marketing stress on fecal excretion of Salmonella spp in feeder calves*. *Am J Vet Res*, 1990. 51(6): p. 866-9.
 111. Hurd, H.S., et al., *Salmonella enterica infections in market swine with and without transport and holding*. *Appl Environ Microbiol*, 2002. 68(5): p. 2376-81.
 112. Hurd, H.S., et al., *The effect of lairage on Salmonella isolation from market swine*. *J Food Prot*, 2001. 64(7): p. 939-44.
 113. Isaacson, R., et al., *Effect of transportation and feed withdrawal on shedding of Salmonella typhimurium among experimentally infected pigs*. *American journal of veterinary research*, 1999. 60 9: p. 1155-8.
 114. Marg, H., et al., *Influence of long-time transportation stress on re-activation of Salmonella typhimurium DT104 in experimentally infected pigs*. *Berl Munch Tierarztl Wochenschr*, 2001. 114(9-10): p. 385-8.
 115. Williams, L.P., Jr. and K.W. Newell, *Salmonella excretion in joy-riding pigs*. *Am J Public Health Nations Health*, 1970. 60(5): p. 926-9.
 116. USDA-APHIS. *Escherichia coli O157 in United States feed-lots*. February 16, 2017]; Available from: <https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/monitoring-and-surveillance/nahms>. Accessed Mar 6, 2023.
 117. Webb, C.R., *Investigating the potential spread of infectious diseases of sheep via agricultural shows in Great Britain*. *Epidemiol Infect*, 2006. 134(1): p. 31-40.
 118. Garber, L.P., et al., *Risk factors for fecal shedding of Escherichia coli O157:H7 in dairy calves*. *J Am Vet Med Assoc*, 1995. 207(1): p. 46-9.
 119. Hancock, D.D., et al., *The prevalence of Escherichia coli O157:H7 in dairy and beef cattle in Washington State*. *Epidemiol Infect*, 1994. 113(2): p. 199-207.
 120. Hancock, D.D., et al., *A longitudinal study of Escherichia coli O157 in fourteen cattle herds*. *Epidemiol Infect*, 1997. 118(2): p. 193-5.
 121. Keen, J.E. and R.O. Elder, *Isolation of shiga-toxigenic Escherichia coli O157 from hide surfaces and the oral cavity of finished beef feedlot cattle*. *J Am Vet Med Assoc*, 2002. 220(6): p. 756-63.
 122. Payne, C.J.I., et al., *Vero Cytotoxin–Producing Escherichia coli O157 Gastroenteritis in Farm Visitors, North Wales*. *Emerging Infectious Disease journal*, 2003. 9(5): p. 526.
 123. Centers for Disease Control and Prevention, *Outbreaks of Escherichia coli O157:H7 infection and cryptosporidiosis associated with drinking unpasteurized apple cider--Connecticut and New York, October 1996*. *MMWR Morb Mortal Wkly Rep*, 1997. 46(1): p. 4-8.
 124. De Schrijver, K., et al., *Outbreak of verocytotoxin-producing E. coli O145 and O26 infections associated with the consumption of ice cream produced at a farm, Belgium, 2007*. *Euro Surveill*, 2008. 13(7).
 125. Djuretic, T., P.G. Wall, and G. Nichols, *General outbreaks of infectious intestinal disease associated with milk and dairy products in England and Wales: 1992 to 1996*. *Commun Dis Rep CDR Rev*, 1997. 7(3): p. R41-5.
 126. Korlath, J.A., et al., *A point-source outbreak of campylobacteriosis associated with consumption of raw milk*. *J Infect Dis*, 1985. 152(3): p. 592-6.
 127. Sharp, J.C., *Infections associated with milk and dairy products in Europe and North America, 1980-85*. *Bull World Health Organ*, 1987. 65(3): p. 397-406.
 128. DeSilva, M.B., et al., *Communitywide cryptosporidiosis outbreak associated with a surface water-supplied municipal water system--Baker City, Oregon, 2013*. *Epidemiol Infect*, 2016. 144(2): p. 274-84.
 129. Bopp, D.J., et al., *Detection, isolation, and molecular subtyping of Escherichia coli O157:H7 and Campylobacter jejuni associated with a large waterborne outbreak*. *J Clin Microbiol*, 2003. 41(1): p. 174-80.
 130. Centers for Disease Control and Prevention, *Outbreak of Escherichia coli O157:H7 and Campylobacter among attendees of the Washington County Fair-New York, 1999*. *MMWR Morb Mortal Wkly Rep*, 1999. 48(36): p. 803-5.
 131. *Waterborne outbreak of gastroenteritis associated with a contaminated municipal water supply, Walkerton, Ontario, May-June 2000*. *Canada communicable disease report = Releve des maladies transmissibles au Canada*, 2000. 26 20: p. 170-3.
 132. Doorduyn, Y., et al., *Risk factors for Salmonella Enteritidis and Typhimurium (DT104 and non-DT104) infections in The Netherlands: predominant roles for raw eggs in Enteritidis and sandboxes in Typhimurium infections*. *Epidemiol Infect*, 2006. 134(3): p. 617-26.
 133. Varma, J.K., et al., *An outbreak of Escherichia coli O157 infection following exposure to a contaminated building*. *JAMA*, 2003. 290(20): p. 2709-12.
 134. Curran K, H.K., Singh T, et al. , *Notes from the field: outbreak of Escherichia coli O157:H7 infections associated with dairy education event attendance—Whatcom County, Washington, 2015*. *MMWR Morb Mortal Wkly Rep* 2015, 2015(56): p. 1285–1287.
 135. Luna, S., et al., *Outbreak of E. coli O157:H7 Infections Associated with Exposure to Animal Manure in a Rural Community - Arizona and Utah, June-July 2017*. *MMWR Morb Mortal Wkly Rep*, 2018. 67(23): p. 659-662.

136. Cavallo, S.J., et al., *Human outbreak of Salmonella Typhimurium associated with exposure to locally made chicken jerky pet treats, New Hampshire, 2013*. Foodborne Pathog Dis, 2015. 12(5): p. 441-6.
137. Behraves, C.B., et al., *Human Salmonella infections linked to contaminated dry dog and cat food, 2006-2008*. Pediatrics, 2010. 126(3): p. 477-83.
138. Centers for Disease Control and Prevention, *Notes from the field: Human Salmonella infantis infections linked to dry dog food--United States and Canada, 2012*. MMWR Morb Mortal Wkly Rep, 2012. 61(23): p. 436.
139. Canada, P.H.A.o., *Human health risk from exposure to natural dog treats*. Can Commun Dis Rep, 2000. 26(6): p. 41-2.
140. Clark, C., et al., *Characterization of Salmonella associated with pig ear dog treats in Canada*. J Clin Microbiol, 2001. 39(11): p. 3962-8.
141. Pitout, J.D., et al., *Association between handling of pet treats and infection with Salmonella enterica serotype newport expressing the AmpC beta-lactamase, CMY-2*. J Clin Microbiol, 2003. 41(10): p. 4578-82.
142. Centers for Disease, C. and Prevention, *Human salmonellosis associated with animal-derived pet treats--United States and Canada, 2005*. MMWR Morb Mortal Wkly Rep, 2006. 55(25): p. 702-5.
143. Khamesipour, F., et al., *A systematic review of human pathogens carried by the housefly (Musca domestica L.)*. BMC Public Health, 2018. 18(1): p. 1049.
144. Conrad, C.C., et al., *Farm Fairs and Petting Zoos: A Review of Animal Contact as a Source of Zoonotic Enteric Disease*. Foodborne Pathog Dis, 2017. 14(2): p. 59-73.
145. McMillian, M., et al., *Risk behaviors for disease transmission among petting zoo attendees*. J Am Vet Med Assoc, 2007. 231(7): p. 1036-8.
146. Koski, L, et al. *Foodborne illness outbreaks linked to unpasteurised milk and relationship to changes in state laws – United States, 1998–2018*. Epidemiology & Infection. 2022; 150:e183
147. Teutsch, S.M., et al., *Epidemic toxoplasmosis associated with infected cats*. N Engl J Med, 1979. 300(13): p. 695-9.
148. Jones, J.L., et al., *Follow-up of the 1977 Georgia Outbreak of Toxoplasmosis*. Am J Trop Med Hyg, 2016. 94(6): p. 1299-300.
149. Simon, J.A., et al., *Spatial distribution of soil contaminated with Toxoplasma gondii oocysts in relation to the distribution and use of domestic cat defecation sites on dairy farms*. Int J Parasitol, 2017. 47(6): p. 357-367.
150. Centers for Disease Control and Prevention, *Outbreak of cutaneous larva migrans at a children's camp--Miami, Florida, 2006*. MMWR Morb Mortal Wkly Rep, 2007. 56(49): p. 1285-7.
151. Glickman, L.T. and P.M. Schantz, *Epidemiology and pathogenesis of zoonotic toxocariasis*. Epidemiol Rev, 1981. 3: p. 230-50.
152. Sorgan, M.H., et al., *A survey of canine toxocariasis and toxocaral soil contamination in Essex County, New Jersey*. Am J Public Health, 1980. 70(11): p. 1207-8.
153. Dada, B.J. and W.D. Lindquist, *Prevalence of Toxocara spp. eggs in some public grounds and highway rest areas in Kansas*. J Helminthol, 1979. 53(2): p. 145-6.
154. Centers for Disease Control and Prevention, *Raccoon roundworms in pet kinkajous--three states, 1999 and 2010*. MMWR Morb Mortal Wkly Rep, 2011. 60(10): p. 302-5.
155. Ryan, U. and A. Zahedi, *Molecular epidemiology of giardiasis from a veterinary perspective*. Adv Parasitol, 2019. 106: p. 209-254.
156. Ryan, U. and S.M. Caccio, *Zoonotic potential of Giardia*. Int J Parasitol, 2013. 43(12-13): p. 943-56.
157. Tsui, C.K., et al., *Beaver Fever: Whole-Genome Characterization of Waterborne Outbreak and Sporadic Isolates To Study the Zoonotic Transmission of Giardiasis*. mSphere, 2018. 3(2).
158. Conners, E.E., et al., *Giardiasis Outbreaks - United States, 2012-2017*. MMWR Morb Mortal Wkly Rep, 2021. 70(9): p. 304-307.
159. KXAN. *Child may have been exposed to rabies at Zilker Park*; Available from: <https://www.kxan.com/news/child-may-have-been-exposed-to-rabies-at-zilker-park/> Accessed Mar 6, 2023.
160. Centers for Disease Control and Prevention, *Mass treatment of humans exposed to rabies--New Hampshire, 1994*. MMWR Morb Mortal Wkly Rep, 1995. 44(26): p. 484-6.
161. Chang, H.G., et al., *Public health impact of reemergence of rabies, New York*. Emerg Infect Dis, 2002. 8(9): p. 909-13.
162. Bender, J.B., et al., *Reports of zoonotic disease outbreaks associated with animal exhibits and availability of recommendations for preventing zoonotic disease transmission from animals to people in such settings*. J Am Vet Med Assoc, 2004. 224(7): p. 1105-9.
163. Centers for Disease Control and Prevention, *Public health response to a potentially rabid bear cub--Iowa, 1999*. MMWR Morb Mortal Wkly Rep, 1999. 48(42): p. 971-3.
164. Centers for Disease Control and Prevention, *Multiple human exposures to a rabid bear cub at a petting zoo and barnwarming--Iowa, August 1999*. MMWR Morb Mortal Wkly Rep, 1999. 48(34): p. 761.
165. Centers for Disease Control and Prevention. *Horse Stabled at Tennessee Walking Horse 2006 National Celebration Tested Positive for Rabies*. 2006 April 4, 2017]; Available from: <https://www.cdc.gov/rabies/resources/news/2006-09-09.html>.

166. Robbins, A., et al., *Bat Incidents at Children's Camps, New York State, 1998–2002*. Emerging Infectious Disease journal, 2005. 11(2): p. 302.
167. Willoughby, R.E., Jr., et al., *Survival after treatment of rabies with induction of coma*. N Engl J Med, 2005. 352(24): p. 2508-14.
168. Centers for Disease Control and Prevention, *Tularemia associated with a hamster bite--Colorado, 2004*. MMWR Morb Mortal Wkly Rep, 2005. 53(51): p. 1202-3.
169. Scheftel, J.M., et al., *Tularaemia in Minnesota: case report and brief epidemiology*. Zoonoses Public Health, 2010. 57(7-8): p. e165-9.
170. Talan, D.A., et al., *Bacteriologic analysis of infected dog and cat bites. Emergency Medicine Animal Bite Infection Study Group*. N Engl J Med, 1999. 340(2): p. 85-92.
171. Cohen, J.I., et al., *Recommendations for prevention of and therapy for exposure to B virus (cercopithecine herpesvirus 1)*. Clin Infect Dis, 2002. 35(10): p. 1191-203.
172. Centers for Disease Control and Prevention, *Fatal Cercopithecine herpesvirus 1 (B virus) infection following a mucocutaneous exposure and interim recommendations for worker protection*. MMWR Morb Mortal Wkly Rep, 1998. 47(49): p. 1073-6, 1083.
173. Hullinger, et al., *Dermatophytosis in show lambs in the United States*. Vet Dermatol, 1999. 10(1): p. 73-76.
174. Lederman, E.R., et al., *ORF virus infection in children: clinical characteristics, transmission, diagnostic methods, and future therapeutics*. Pediatr Infect Dis J, 2007. 26(8): p. 740-4.
175. Kurth, A., et al., *Rat-to-elephant-to-human transmission of cowpox virus*. Emerg Infect Dis, 2008. 14(4): p. 670-1.
176. Ninove, L., et al., *Cowpox virus transmission from pet rats to humans, France*. Emerg Infect Dis, 2009. 15(5): p. 781-4.
177. Kile, J.C., et al., *Transmission of monkeypox among persons exposed to infected prairie dogs in Indiana in 2003*. Arch Pediatr Adolesc Med, 2005. 159(11): p. 1022-5.
178. Centers for Disease Control and Prevention, *Update: multistate outbreak of monkeypox--Illinois, Indiana, Kansas, Missouri, Ohio, and Wisconsin, 2003*. MMWR Morb Mortal Wkly Rep, 2003. 52(27): p. 642-6.
179. Nemetz T, S.E.J., *Zoonotic diseases*, in *Fish Medicine*. 1990, WB Saunders Co: Philadelphia. p. 214-220.
180. Gray, S.F., et al., *Fish tank granuloma*. BMJ, 1990. 300(6731): p. 1069-70.
181. Lewis, F.M., B.J. Marsh, and C.F. von Reyn, *Fish tank exposure and cutaneous infections due to Mycobacterium marinum: tuberculin skin testing, treatment, and prevention*. Clin Infect Dis, 2003. 37(3): p. 390-7.
182. Lucky, A.W., et al., *Avian mite bites acquired from a new source--pet gerbils: report of 2 cases and review of the literature*. Arch Dermatol, 2001. 137(2): p. 167-70.
183. Angarano, D.W. and L.C. Parish, *Comparative dermatology: parasitic disorders*. Clin Dermatol, 1994. 12(4): p. 543-50.
184. Arlian, L.G., *Biology, host relations, and epidemiology of Sarcoptes scabiei*. Annu Rev Entomol, 1989. 34: p. 139-61.
185. Scott, D.W. and R.T. Horn, Jr., *Zoonotic dermatoses of dogs and cats*. Vet Clin North Am Small Anim Pract, 1987. 17(1): p. 117-44.
186. Barrington, G.M., J.M. Gay, and J.F. Evermann, *Biosecurity for neonatal gastrointestinal diseases*. Vet Clin North Am Food Anim Pract, 2002. 18(1): p. 7-34.
187. Centers for Disease Control and Prevention, *Influenza A (H3N2) variant virus*. Available from: <https://www.cdc.gov/flu/swineflu/variant/h3n2v-cases.htm>. Accessed Mar 6, 2023.
188. Bowman, A.S., et al., *Subclinical influenza virus A infections in pigs exhibited at agricultural fairs, Ohio, USA, 2009-2011*. Emerg Infect Dis, 2012. 18(12): p. 1945-50.
189. Yassine, H.M., et al., *Characterization of triple reassortant H1N1 influenza A viruses from swine in Ohio*. Vet Microbiol, 2009. 139(1-2): p. 132-9.
190. Centers for Disease Control and Prevention, *Swine-origin influenza A (H3N2) virus infection in two children--Indiana and Pennsylvania, July-August 2011*. MMWR Morb Mortal Wkly Rep, 2011. 60(35): p. 1213-5.
191. Centers for Disease Control and Prevention, *Limited human-to-human transmission of novel influenza A (H3N2) virus--Iowa, November 2011*. MMWR Morb Mortal Wkly Rep, 2011. 60(47): p. 1615-7.
192. Centers for Disease Control and Prevention, *Influenza A (H3N2) variant virus-related hospitalizations: Ohio, 2012*. MMWR Morb Mortal Wkly Rep, 2012. 61: p. 764-7.
193. Centers for Disease Control and Prevention *Update: Influenza A (H3N2)v transmission and guidelines - five states, 2011*. MMWR Morb Mortal Wkly Rep, 2012. 60(51-52): p. 1741-4.
194. Wong, K.K., et al., *Outbreak of influenza A (H3N2) variant virus infection among attendees of an agricultural fair, Pennsylvania, USA, 2011*. Emerg Infect Dis, 2012. 18(12): p. 1937-44.
195. National Association of State Public Health Veterinarians, *Measures to Minimize Influenza Transmission at Swine Exhibitions, 2018*. October 31, 2022]; Available from: <http://nasphv.org/Documents/InfluenzaTransmissionAtSwineExhibitions2018.pdf>.
196. Vincent, A.L., et al., *Characterization of an influenza A virus isolated from pigs during an outbreak of respiratory disease in swine and people during a county fair in the United States*. Vet Microbiol, 2009. 137(1-2): p. 51-9.
197. Wells, D.L., et al., *Swine influenza virus infections. Transmission from ill pigs to humans at a Wisconsin agricultural fair*

- and subsequent probable person-to-person transmission. JAMA, 1991. 265(4): p. 478-81.
198. Cox, C.M., et al., *Swine influenza virus A (H3N2) infection in human, Kansas, USA, 2009*. Emerg Infect Dis, 2011. 17(6): p. 1143-4.
 199. Centers for Disease Control and Prevention. *Avian influenza A (H7N2) in cats in animal shelters in NY; one human infection*. Available from: <https://www.cdc.gov/flu/spotlights/avian-influenza-cats.htm>. Accessed Mar 6, 2023.
 200. Karasin, A.I., S. Carman, and C.W. Olsen, *Identification of human H1N2 and human-swine reassortant H1N2 and H1N1 influenza A viruses among pigs in Ontario, Canada (2003 to 2005)*. J Clin Microbiol, 2006. 44(3): p. 1123-6.
 201. Vincent, A.L., et al., *Swine influenza viruses a North American perspective*. Adv Virus Res, 2008. 72: p. 127-54.
 202. Zhou, N.N., et al., *Genetic reassortment of avian, swine, and human influenza A viruses in American pigs*. J Virol, 1999. 73(10): p. 8851-6.
 203. Van Reeth, K., I. Brown and C.W. Olsen, *Influenza Virus*, in J.J. Zimmerman, L.A. Karriker, S. A. Ramirez, K.J. Schwartz and G.W. Stevenson (eds.) *Diseases of Swine, 10th Edition*. 2012, John Wiley and Son, Inc: Ames, IA.
 204. Murphree, R., et al., *Elephant-to-human transmission of tuberculosis, 2009*. Emerg Infect Dis, 2011. 17(3): p. 366-71.
 205. Kiers, A., et al., *Transmission of Mycobacterium pinnipedii to humans in a zoo with marine mammals*. Int J Tuberc Lung Dis, 2008. 12(12): p. 1469-73.
 206. Oh, P., et al., *Human exposure following Mycobacterium tuberculosis infection of multiple animal species in a Metropolitan Zoo*. Emerg Infect Dis, 2002. 8(11): p. 1290-3.
 207. Michalak, K., et al., *Mycobacterium tuberculosis infection as a zoonotic disease: transmission between humans and elephants*. Emerg Infect Dis, 1998. 4(2): p. 283-7.
 208. Zlot, A., et al., *Diagnosis of Tuberculosis in Three Zoo Elephants and a Human Contact - Oregon, 2013*. MMWR Morb Mortal Wkly Rep, 2016. 64(52): p. 1398-402.
 209. Sub-committee, U.A.H.A.E.T. *Guidelines for the control of tuberculosis in elephants 2012*. 2012; Available from: <https://www.usaha.org/upload/Committee/Tuberculosis/TB%20Guidelines%202012%20Draft%20revision%2020April2012.pdf>. Accessed Mar 6, 2023.
 210. Whelan, J., et al., *Visits on 'lamb-viewing days' at a sheep farm open to the public was a risk factor for Q fever in 2009*. Epidemiol Infect, 2012. 140(5): p. 858-64.
 211. McQuiston, J.H. and J.E. Childs, *Q fever in humans and animals in the United States*. Vector Borne Zoonotic Dis, 2002. 2(3): p. 179-91.
 212. Anderson, A., et al., *Diagnosis and management of Q fever--United States, 2013: recommendations from CDC and the Q Fever Working Group*. MMWR Recomm Rep, 2013. 62(RR-03): p. 1-30.
 213. Balsamo, G., et al., *Compendium of Measures to Control Chlamydia psittaci Infection Among Humans (Psittacosis) and Pet Birds (Avian Chlamydiosis), 2017*. J Avian Med Surg, 2017. 31(3): p. 262-282.
 214. Christensen, A.L., J.O. Jarløv, and S. Ingeberg, *[The risk of ornithosis among the staff of Copenhagen Zoo]*. Ugeskrift for laeger, 1990. 152(12): p. 818-820.
 215. Schlossberg, D., et al., *An epidemic of avian and human psittacosis*. Arch Intern Med, 1993. 153(22): p. 2594-6.
 216. Eidson, M., *Psittacosis/avian chlamydiosis*. J Am Vet Med Assoc, 2002. 221(12): p. 1710-2.
 217. Hyde, S.R. and K. Benirschke, *Gestational psittacosis: case report and literature review*. Mod Pathol, 1997. 10(6): p. 602-7.
 218. Gherman, R.B., L.L. Leventis, and R.C. Miller, *Chlamydial psittacosis during pregnancy: a case report*. Obstet Gynecol, 1995. 86(4 Pt 2): p. 648-50.
 219. Khatib, R., et al., *Severe psittacosis during pregnancy and suppression of antibody response with early therapy*. Scand J Infect Dis, 1995. 27(5): p. 519-21.
 220. Centers for Disease Control and Prevention, *Outbreaks of Escherichia coli O157:H7 infections among children associated with farm visits--Pennsylvania and Washington, 2000*. MMWR Morb Mortal Wkly Rep, 2001. 50(15): p. 293-7.
 221. Anderson, M.E. and J.S. Weese, *Video observation of hand hygiene practices at a petting zoo and the impact of hand hygiene interventions*. Epidemiol Infect, 2012. 140(1): p. 182-90.
 222. Erdozain, G., et al., *Observation of public health risk behaviours, risk communication and hand hygiene at Kansas and Missouri petting zoos--2010-2011*. Zoonoses Public Health, 2013. 60(4): p. 304-10.
 223. Centers for Disease Control and Prevention. *Handwashing: clean hands save lives. Show me the science-when & how to use hand sanitizer*. Available from: www.cdc.gov/handwashing/show-me-the-science-hand-sanitizer.html. Accessed Mar 6, 2023.
 224. Keen, J.E., L.M. Durso, and T.P. Meehan, *Isolation of Salmonella enterica and Shiga-toxigenic Escherichia coli O157 from feces of animals in public contact areas of United States zoological parks*. Appl Environ Microbiol, 2007. 73(1): p. 362-5.
 225. Crump, J.A., et al., *Outbreaks of Escherichia coli O157 infections at multiple county agricultural fairs: a hazard of mixing cattle, concession stands and children*. Epidemiol Infect, 2003. 131(3): p. 1055-62.
 226. Hoek, M.R., et al., *Outbreak of Cryptosporidium parvum among children after a school excursion to an adventure farm, south west England*. J Water Health, 2008. 6(3): p. 333-8.

227. Hedicán, E., et al., *Salmonellosis outbreak due to chicken contact leading to a foodborne outbreak associated with infected delicatessen workers*. Foodborne Pathog Dis, 2010. 7(8): p. 995-7.
228. Centers for Disease Control and Prevention, *Outbreak of Shiga toxin-producing Escherichia coli O157 infection associated with a day camp petting zoo--Pinellas County, Florida, May-June 2007*. MMWR Morb Mortal Wkly Rep, 2009. 58(16): p. 426-8.
229. Smith, K.E., et al., *Outbreaks of salmonellosis at elementary schools associated with dissection of owl pellets*. Vector borne and zoonotic diseases (Larchmont, N.Y.), 2005. 5(2): p. 133-136.
230. American Veterinary Medical Association Task Force on Canine, A. and I. Human-Canine, *A community approach to dog bite prevention*. J Am Vet Med Assoc, 2001. 218(11): p. 1732-49.
231. Canada, P.H.A. *Injuries associated with... equestrian activities*. CHIRPP database, summary data for 1996, all ages. Available from: <https://www.canada.ca/en/public-health/services/injury-prevention/canadian-hospitals-injury-reporting-prevention-program/injury-reports/injuries-associated-equestrian-activities.html>. Accessed Mar 6, 2023.
232. Associated Press. *Teen killed by tiger at Kansas sanctuary*. Available from: <https://www.foxnews.com/story/teen-killed-by-tiger-at-kansas-sanctuary>. Accessed Mar 6, 2023.
233. KNKX. *Llama attacks woman at B.C. petting zoo*. 2011. Available from: <https://www.knkn.org/other-news/2011-08-04/llama-viciously-attacks-elderly-woman-at-b-c-petting-zoo>. Accessed Mar 6, 2023.
234. Cherry, C., et al., *Notes from the Field: Injuries Associated with Bison Encounters - Yellowstone National Park, 2015*. MMWR Morb Mortal Wkly Rep, 2016. 65(11): p. 293-4.
235. KLTV. *Two boys lucky to be okay after being bitten by water moccasin in class*. 2008. Available from: www.kltv.com/story/9216055/two-boys-lucky-to-be-okay-after-being-bitten-by-water-moccasin-in-class. Accessed Mar 6, 2023.
236. Associated Press. *Ten-year-old bitten by camel at Virginia safari park gets \$155,000 settlement*. 2016. Available from: www.theguardian.com/us-news/2016/aug/20/virginia-safari-park-camel-bite-lawsuit-settlement. Accessed Mar 6, 2023.
237. Kahn, A., et al., *Child victims of dog bites treated in emergency departments: a prospective survey*. Eur J Pediatr, 2003. 162(4): p. 254-8.
238. Casemore, D. *Educational farm visits and associated infection hazards*. Commun Dis Rep CDR Rev, 1989. 19:3.
239. Dawson, A., et al., *Farm visits and zoonoses*. Commun Dis Rep CDR Rev, 1995. 5(6): p. R81-6.
240. Warshawsky B, H.B., Gutmanis I. *An E. coli O157:H7 outbreak associated with an animal exhibit: Middlesex-London Health Unit Investigation and recommendations—executive summary*. 1999. Available from: www.healthunit.com/uploads/mlhu-e-coli-o157h7-outbreak.pdf. Accessed Mar 6, 2023.
241. Hoss, A., et al., *State Laws Requiring Hand Sanitation Stations at Animal Contact Exhibits-United States, March-April 2016*. MMWR Morb Mortal Wkly Rep, 2017. 66(1): p. 16-18.
242. *North Carolina Office of Administrative Hearings*. Available from: <http://reports.oah.state.nc.us/ncac/title%2002%20-%20agriculture%20and%20consumer%20services/chapter%2052%20-%20veterinary/subchapter%20k/subchapter%20k%20rules.pdf>. Accessed Mar 6, 2023.
243. AZA. *The accreditation standards and related policies: 2023 edition*. Available from: www.aza.org/assets/2332/aza-accreditation-standards.pdf. Accessed Mar 6, 2023.
244. USDA-APHIS. *Licensing and registration under the Animal Welfare Act*. Available from: <https://www.aphis.usda.gov/aphis/ourfocus/animalwelfare/apply/licensing-and-registration-application-packets>. Accessed Mar 6, 2013.
245. Centers for Disease Control and Prevention. *Communication Materials: Salmonella and Enteric Zoonoses*. Available from: <https://www.cdc.gov/salmonella/reportspubs/communication-materials.html>. Accessed Mar 6, 2023.
246. National Association of State Public Health Veterinarians, *Compendium of Animal Rabies Prevention and Control, 2016*. J Am Vet Med Assoc, 2016. 248(5): p. 505-17.
247. National Association of State Animal Health Officials, et. al. *Prevention and control of Coxiella burnetii infection among humans and animals: guidance for a coordinated public health and animal health response, 2013*. Available from: www.nasphv.org/Documents/Q_Fever_2013.pdf. Accessed Mar 6, 2023.
248. National Association of State Public Health Veterinarians. *Zoonotic influenza*. Available from: www.nasphv.org/documentsCompendiaZoonoticInfluenza.html. Accessed Mar 6, 2023.
249. Bondeson, L., *Assessment of measures to prevent disease associated with animals in agricultural fairs--Maine, 2008*. Am J Infect Control, 2009. 37(8): p. 665-7.
250. National Association of State Public Health Veterinarians. *Animals in public settings toolkit*. Available from: www.nasphv.org/documentsCompendiumAnimals.html. Accessed Mar 6, 2023.
251. Nichols, M., et al., *Outbreak of Multidrug-Resistant Salmonella Heidelberg Infections Linked to Dairy Calf Exposure, United States, 2015-2018*. Foodborne Pathog Dis, 2022. 19(3): p. 199-208.
252. Centers for Disease Control and Prevention. *Hygiene Practices Around Animals*. Available from: <https://www.cdc.gov/healthypets/index.html>. Accessed Mar 6, 2023.

253. Michaels, B., et al., *Water temperature as a factor in handwashing efficacy*. Food Service Technology, 2002. 2(3): p. 139-149.
254. Messenger, A.M., A.N. Barnes, and G.C. Gray, *Reverse zoonotic disease transmission (zooanthroponosis): a systematic review of seldom-documented human biological threats to animals*. PLoS One, 2014. 9(2): p. e89055.
255. Center for Food Security and Public Health. *Disinfection*. Available from: www.cfsph.iastate.edu/Disinfection/index.php. Accessed Mar 6, 2023.
256. Centers for Disease Control and Prevention. *Raw Milk*. April 24, 2021]; Available from: <https://www.cdc.gov/foodsafety/rawmilk/raw-milk-index.html>.
257. Virginia Department of Public Health. *Non-human primates*. Available from: www.vdh.virginia.gov/environmental-epidemiology/. Accessed Mar 6, 2023.
258. Swift, J.M., et al., *Efficacy of an Escherichia coli O157:H7 SRP Vaccine in Orally Challenged Goats and Strain Persistence Over Time*. Foodborne Pathog Dis, 2017. 14(3): p. 160-166.
259. American Association of Zoo Veterinarians. *American Association of Zoo Veterinarians Infectious Disease Committee manual 2013: tuberculosis in non-human primates*. Available from: <https://www.aazv.org/page/754>. Accessed Mar 6, 2023.
260. Centers for Disease Control and Prevention. *Q fever safety at livestock birthing exhibits: information for operators and managers*. February 14, 2017]; Available from: www.cdc.gov/qfever/pdfs/qfever-factsheet.pdf.
261. McBride, D.S., et al., *Shortening Duration of Swine Exhibitions to Reduce Risk for Zoonotic Transmission of Influenza A Virus*. Emerg Infect Dis, 2022. 28(10): p. 2035-2042.
262. Bowman, A.S., et al., *Influenza A(H3N2) Virus in Swine at Agricultural Fairs and Transmission to Humans, Michigan and Ohio, USA, 2016*. Emerg Infect Dis, 2017. 23(9): p. 1551-1555.
263. Yassine, H.M., C.W. Lee, and Y.M. Saif, *Interspecies transmission of influenza A viruses between Swine and poultry*. Curr Top Microbiol Immunol, 2013. 370: p. 227-40.

Appendix 1

Selected Zoonotic Diseases of Importance in Public Settings in the United States, 2023.

Disease	Agent	Most common species associated with transmission to humans	Transmission route to humans	Symptoms in humans
Acariasis (mite infestation)	<i>Sarcoptes scabiei</i> (species-specific variants), <i>Notoedres cati</i> , <i>Cheyletiella</i> spp, other species of mites	Dogs, cats, horses, goats, sheep, swine, birds, rats, rabbits	Direct or indirect contact*	Itchy skin lesions
B virus infection	B virus	Macaque monkeys	Bites, scratches	Localized skin lesions, influenza-like symptoms, brain inflammation
Bartonellosis (cat scratch disease)	<i>Bartonella henselae</i> , other <i>Bartonella</i> spp	Cats (flea dirt)	Scratches, bites	Fever, malaise, enlarged/painful lymph nodes, skin lesions at site of scratch or bite
Baylisascariasis	<i>Baylisascaris procyonis</i>	Raccoons, occasionally dogs	Fecal-oral†	Variable depending on system the parasite moves through (eye, organs, brain)
Brucellosis	<i>Brucella</i> spp	Dogs, cervids, feral swine, bison, marine mammals	Ingestion, inhalation, contact with mucous membranes	Variable, nonspecific febrile illness
Campylobacteriosis	<i>Campylobacter jejuni</i> , other <i>Campylobacter</i> spp	Poultry, cattle, sheep, goats, swine, dogs, cats, turtles	Fecal-oral contact	Gastroenteritis, fever; usually self-limiting
Chlamydiosis	<i>Chlamydophila abortus</i> , <i>Chlamydophila felis</i>	Sheep, goats, llamas, cats, cattle	Inhalation, fecal-oral contact	Miscarriage, septicemia
Contagious ecthyma (orf)	Parapoxvirus	Sheep, goats (sore mouth)	Direct or indirect contact	Skin papules, enlarged/painful lymph nodes, influenza-like illness
Cowpox infection	Cowpox virus	Cattle, rodents, cats, exotic animals	Direct contact	Skin pustules, febrile illness
Cryptosporidiosis	<i>Cryptosporidium parvum</i>	Cattle (typically calves), sheep, goats	Fecal-oral contact	Gastroenteritis
Cutaneous larva (hookworm)	<i>Ancylostoma braziliense</i> , <i>Ancylostoma caninum</i>	Dogs, cats	Direct contact with contaminated soil	Larva migrans skin lesions
Dermatophytosis (ringworm)	<i>Microsporum</i> spp, <i>Trichophyton</i> spp, <i>Epidermophyton</i> spp	Cats, dogs, cattle, goats, sheep, horses, rabbits, rodents, hedgehogs	Direct or indirect contact	Most commonly a ring-shaped rash
Influenza	Influenza A virus	Swine, poultry	Inhalation, direct or indirect contact	Fever, malaise, muscle and joint pain
Leptospirosis	<i>Leptospira</i> spp	Swine, cattle, dogs, rodents	Direct or indirect contact, inhalation	Fever, headache, muscle aches, kidney/liver failure is possible
Listeriosis	<i>Listeria monocytogenes</i>	Cattle, sheep, goats, pigs, dogs, cats	Fecal-oral contact	Fever, headaches, muscle ache, stiff neck, confusion, seizures, miscarriage
Mpox (Monkeypox)	Monkeypox virus	Rodents, non-human primates	Direct or indirect contact, bites, inhalation	Influenza-like symptoms followed by skin lesions
Mycobacteriosis (nontuberculous)	<i>Mycobacterium marinum</i>	Aquarium fish	Direct contact with infected fish or contaminated water, aerosol	Skin lesions

Pasteurellosis	<i>Pasteurella multocida</i> and other species	Dogs, cats, rabbits	Bites, scratches, contact with mucous membranes	Wound infections
Psittacosis	<i>Chlamydia psittaci</i>	Pet birds, poultry	Inhalation, bites	Fever, headache, muscle aches, cough
Q fever	<i>Coxiella burnetii</i>	Goats, sheep, cattle	Inhalation	Influenza-like symptoms, pneumonia
Rabies	Rabies virus	Domestic and wild mammals	Bites, saliva contact with mucous membranes or wounds	Acute, progressive, fatal neurologic disease
Rat bite fever	<i>Streptobacillus moniliformis</i> , <i>Spirillum minus</i>	Rats, mice, gerbils	Bites, scratches	Fever, severe muscle and joint pain
Salmonellosis	<i>Salmonella</i> spp	Reptiles, amphibians, poultry, swine, cattle, goats, horses, rodents, hedgehogs	Fecal-oral contact	Gastroenteritis
Staphylococcosis	<i>Staphylococcus</i> spp	Swine, dogs, cats	Bites, scratches	Localized skin and soft tissue infections
STEC infection	Shiga toxin-producing <i>Escherichia. coli</i>	Cattle, goats, sheep, deer	Fecal-oral contact	Gastroenteritis, hemolytic-uremic syndrome
Streptococcosis	<i>Streptococcus</i> spp	Swine, dogs, cats	Bites, scratches	Localized skin and soft tissue infections
Toxoplasmosis	<i>Toxoplasma gondii</i>	Cats	Fecal-oral contact	Enlarged/painful lymph nodes, mild influenza-like symptoms
Tuberculosis	<i>Mycobacterium tuberculosis</i> complex	Elephants, cattle, nonhuman primates	Inhalation	Respiratory disease
Visceral larva migrans	<i>Toxocara canis</i> , <i>Toxocara cati</i> , <i>Baylisascaris procyonis</i>	Dogs, cats, raccoons	Fecal-oral contact	Various and nonspecific signs (e.g., fever, lethargy, cough)

* Direct transmission occurs after direct physical contact with the animal, which includes touching, holding, kissing, being bitten, licked, and scratched. Indirect transmission occurs indirectly from an animal through contact with a surface contaminated by the animal's saliva, blood, urine, respiratory secretions, feces, or other bodily fluids or via contaminated human food.

† Fecal – oral transmission occurs when small amounts of feces (possibly not visible) that contain germs are accidentally ingested, often by hand-to-mouth contact.

(Adapted from NASPHV Animal Contact Compendium Committee 2017. "Compendium of Measures to Prevent Disease Associated with Animals in Public Settings, 2017." *J Am Vet Med Assoc* **251**(11): 1268-1292. Reprinted with permission.)

Appendix 2

Animals in Public Settings: Recommendations for Venue Operators, Staff, and Volunteers^[1-4]

All individuals involved with animal contact activity in any public setting should be aware of the following risks for disease and injury associated with animals in public settings:

- Disease and injuries have occurred following contact with animals and their environment.
- Animals that appear healthy can carry harmful germs that can make visitors sick.
- Visitors can pick up harmful pathogens when they touch animals or animal droppings or enter animal environments (even without directly contacting the animals).
- Visitors can decrease the risk from most germs by washing their hands immediately after leaving an animal area. Visitors should wash their hands even if they did not directly contact the animals.
- Visitors can also reduce their exposure to germs by removing or changing clothes and shoes that have been worn in animal environments.
- The risk for developing serious or life-threatening zoonotic disease from contact with animals is higher for some visitors than others, especially children < 5 years of age, persons > 65 years of age, pregnant women, and people with weakened immune systems.
- Direct contact with some animals is inappropriate for some, or all, audiences in public settings.
 - No visitors should have contact with ill animals.
 - Direct contact with preweaned calves, reptiles, amphibians, and live poultry is not appropriate for people at high risk for zoonotic disease transmission, and direct contact with young ruminants of other species (e.g., goats and sheep) is of increased concern for these individuals.
 - Dangerous animals (e.g., nonhuman primates, certain carnivores, other rabies reservoir species, and venomous reptiles) should be prohibited from having direct contact with the public.
 - Live animals, especially reptiles, amphibians, and live poultry, should not be given as prizes at fairs, carnivals, or other events.

Operators and all individuals involved with the animal contact activity should educate visitors (with simple instructions in multiple age-appropriate and language-appropriate formats) about the following before they enter animal areas:

- Risks for disease and injury, including the information that children < 5 years of age, people > 65 years of age, pregnant women, and those with weakened immune systems are at greater risk than others of developing serious zoonotic diseases.
- Appropriate (or thorough) handwashing and assisting children with handwashing immediately after visiting an animal area.
- Avoiding eating, drinking, or placing things in their mouths after animal contact or after visiting an animal area, until they have washed their hands.
- Closely supervising children.
- Awareness that objects such as clothing, shoes, and stroller wheels can become soiled and serve as a source of germs after leaving an animal area.

Operators and all individuals involved with the animal contact activity should take the following steps to maintain a safe environment when animals are present in public settings:

- Design the venue with safety in mind by having designated animal areas, nonanimal areas, and transition areas; temporary exhibits and animal interaction areas used in farm visits, agritourism venues, etc. may need additional measures to minimize risks of injury or disease transmission.
- Do not permit animals other than service animals in nonanimal areas.
- Assign trained staff members to monitor animal contact areas to ensure visitor safety and provide education about risks and best prevention methods.
- Exclude food and beverages, toys, pacifiers, spill-proof cups, baby bottles, and smoking and related activities from animal contact areas.
- Keep the animal areas as clean and disinfected as possible, and limit visitor contact with manure and animal bedding.
- Allow feeding of animals only if contact with animals can be controlled (e.g., over a barrier), and do not provide feed in containers that might be consumed by persons (e.g., ice cream cones).
- Design transition areas for entering and exiting animal areas with appropriate signs or notifications regarding risks associated with animal contact and location of handwashing facilities.
- Maintain handwashing stations that are accessible to children and people with disabilities, and direct visitors to wash their hands immediately upon exiting animal areas.
- Position handwashing stations in places that encourage handwashing when exiting animal areas.
- Maintain handwashing facilities and stations appropriately by conducting routine cleaning and restocking to ensure an adequate supply of paper towels and soap.
- Ensure animals receive appropriate preventive care, including vaccinations and parasite control appropriate for the species.
- Provide potable water for animals.
- Provide handwashing facilities where food and beverages are stored, prepared, served, or consumed.
- Prohibit consumption of unpasteurized dairy products (e.g., raw milk), ciders, and juices.
- Minimize use of animal areas at other times for public activities (e.g., weddings, dances, and barbecues) because germs that can make people sick can persist in the environment.

Handwashing is the most important prevention step for reducing disease transmission associated with animals in public settings.

(Adapted from NASPHV Animal Contact Compendium Committee 2017. "Compendium of Measures to Prevent Disease Associated with Animals in Public Settings, 2017." *J Am Vet Med Assoc* 251(11): 1268-1292. Reprinted with permission.)

Appendix 3

Handwashing Recommendations to Reduce Disease Transmission from Animals in Public Settings^[1-4]

Handwashing is the most important prevention step for reducing disease transmission associated with animals in public settings. Hands should always be washed immediately upon exiting animal areas, even if direct contact with animals did not occur. Handwashing is also important before eating, drinking, or handling food and after removing soiled clothing or shoes.

Correct Handwashing Procedure

- Wet both hands with clean (warm or cold) running water and apply soap.
- Rub hands together for at least 20 seconds (sing "Happy Birthday" twice) to make a lather and be sure to scrub the backs of hands, wrists, between fingers, and under nails.
- Rinse hands thoroughly under running water. Where available, use a disposable paper towel to turn off faucets.
- Dry hands with a clean paper towel or air-dry them. Dispose of used paper towels in a designated receptacle. Do not dry hands on clothing.
- Assist young children with washing, rinsing and drying their hands.

Establishment and Maintenance of Handwashing Facilities or Stations

- Venue staff should encourage visitors to wash hands immediately upon exiting animal areas.
- The number of handwashing stations should be sufficient for the maximum anticipated attendance; facilities should be accessible to children (i.e., low enough for children to reach or equipped with a stool), people with disabilities as well as the general public.
- Handwashing facilities should be conveniently located in transition areas between animal and non-animal areas and in all food concession areas.
- Maintenance of handwashing stations should include routine cleaning and regular restocking to ensure an adequate supply of paper towels and soap.
- Running water should be of sufficient volume and pressure to remove soil from hands. Volume and pressure might be substantially reduced if the water supply is furnished from a holding tank; therefore, a permanent, pressurized water supply is preferable.
- Handwashing stations should be designed so that both hands are free to employ correct handwashing procedures by having automated sensor faucets, water that stays on after hand faucets are turned, or stations that operate using a foot pedal.
- Liquid soap dispensed by a hand pump, foot pump or automatic dispenser is recommended.
- To increase compliance, water temperature should be set at what is considered comfortable.²³⁷

Temporary handwashing stations

- Handwashing stations should be equipped with potable water in a clean enclosed container, liquid hand soap, a greywater container to catch used water, single use paper towels and a trash container. (**Figure 4**)
- A number of cost-effective premade or simple build-your-own designs are available
- Communal basins, in which water is used by more than one person at a time, are not adequate handwashing facilities.

Handwashing Sign Recommendations


- Anywhere human-animal contact occurs in public, signs regarding proper handwashing practices are critical to reduce disease transmission.
- Signs to remind visitors to wash hands should be posted at exits from animal areas, near portable toilets, and in non-animal areas where food is served and consumed.
- Signs outlining proper handwashing instructions should be posted at handwashing stations and in restrooms to encourage proper practices.
- Handwashing signs should be available in multiple age-appropriate and language-appropriate formats.

Recommendations Regarding Hand-Sanitizing Agents

- Washing hands with soap and water is the best way to reduce the number of germs on them. If soap and water are not available, use an alcohol-based hand sanitizer that contains at least 60% alcohol in the interim until hands can be properly washed.
- Visible contamination and dirt should be removed before using hand sanitizers. Hand sanitizers may not be as effective when hands are visibly dirty or greasy.
- Even when hand sanitizer is used, visitors should always wash hands with soap and water as soon as possible after exiting animal areas; alcohol-based hand sanitizers can quickly reduce the number of germs on hands in some situations, but these products are not effective against all germs.

Correct Use of Hand Sanitizers

- Apply the gel to the palm of one hand using the amount suggested on the product label, rub both hands together, and continue to rub the product over all surfaces of your hands and fingers until your hands are dry (about 20 seconds).



Appendix 4

Guidelines for Exhibition of Animals in School and Childcare Settings^{1-4]}

General Recommendations

- Animals are effective and valuable teaching aids, but safeguards are required to reduce the risk for infection and injury. Other entities have developed recommendations similar to those provided here.
- Ensure that teachers and staff know which animal species are inappropriate for their facility and which animals should not be in direct contact with children (See animal-specific recommendations in this Appendix).
- Ensure that personnel providing animals for educational purposes are knowledgeable regarding animal handling and zoonotic disease issues. People or facilities that display animals to the public should be licensed by the USDA.
- Inform parents of the presence of resident or visiting animals as well as the benefits and potential risks associated with animals in schools and childcare settings. Consult with parents to determine special considerations for children who are immunocompromised, have allergies or asthma.
- Educate children about harmful germs that can spread between animals and people and about proper handwashing technique.
- Wash hands immediately after contact with animals, animal products, or feed or after being around animal environments.
- Supervise human-animal contact, particularly involving children < 5 years of age.
- Display animals in enclosed cages or using appropriate restraints.
- Do not allow resident or visiting animals in schools or childcares to roam or fly free or potentially have contact with wild animals.
- Designate specific areas for animal contact to occur. Do not allow food or drink in animal contact areas; do not allow animals in areas where food and drink are stored, prepared, served, or consumed.
- Clean and disinfect all areas where animals and animal products have been present. Children should not perform this task except under adult supervision.
- Do not clean animal cages or enclosures, animal bowls, toys or animal enrichment objects in sinks or other areas used to store, prepare, serve, or consume food and drinks. Designate a utility or laundry sink for cleaning animal cages and other animal contact items.
- Obtain a certificate of veterinary inspection, proof of rabies vaccination, or both according to local and state requirements for the animal species being exhibited. Ensure veterinary care, including preventive health programs for fleas, ticks, mites, and internal parasites is provided as appropriate for the species.

Animal-Specific Recommendations

Do not keep the following animals in facilities with children < 5 years of age and do not allow this age group to have direct contact with these animals as these animals pose a high risk for zoonotic disease transmission or bites:

- Reptiles (e.g., turtles, snakes, and lizards)
- Amphibians (e.g., frogs, toads, salamanders, and newts)
- Live poultry (e.g., chicks, ducklings, and goslings)
- Ferrets: prevent direct contact with these animals to avoid bites.

Refer to the general guidelines regarding species for which specific recommendations are not provided in this section (e.g., nonsittacine birds and domestic dogs, cats, rabbits, and rodents [including mice, rats, hamsters, gerbils, guinea pigs, and chinchillas]).

- Guide dogs, hearing assistance, or other service animals and trained animals used in law enforcement may be present in accordance with recommendations from the sponsoring organizations when they are under the control of a person familiar with the specific animal.
- Psittacine birds (e.g., parrots, parakeets, and cockatiels): Consult the psittacosis compendium¹⁹⁸ and seek veterinary advice.
- Fish: Children < 5 years of age and people with impaired immune systems should not clean aquariums. Wash hands before and after cleaning aquariums, and wear gloves if hands have cuts or wounds or when working with rough rocks or spiny fish. Do not dispose of aquarium water in sinks used for food preparation or for obtaining drinking water.
- Animal products: Assume that products such as owl pellets and frozen rodents used to feed reptiles are contaminated with *Salmonella* organisms. Dissection of owl pellets should not be performed in areas where food is stored, prepared, served, or consumed. Children < 5 years of age should not be allowed to have direct contact with animal products unless the product has been treated to eliminate germs.

Animals Not Recommended in Any Setting Including School or Childcare Settings

- Inherently dangerous or wild animals (e.g., lions, tigers, cougars, and bears).
- Nonhuman primates (e.g., monkeys and apes).
- Mammals that pose a high risk for transmitting rabies (e.g., bats, raccoons, skunks, foxes, and coyotes).
- Aggressive or unpredictable wild or domestic animals.
- Stray animals with unknown health and vaccination history.
- Venomous or toxin-producing spiders, insects, reptiles, and amphibians.
- Animals that pose a high risk for bites (e.g., ferrets). If present in public settings, ferrets should be up-to-date for rabies vaccination.
- Farm animals (e.g., calves, goats, and sheep) should not be displayed to older children in school settings unless meticulous attention to personal hygiene can be ensured because these animals intermittently shed substantial amounts of germs

(Adapted from NASPHV Animal Contact Compendium Committee 2017. "Compendium of Measures to Prevent Disease Associated with Animals in Public Settings, 2017." *J Am Vet Med Assoc* 251(11): 1268-1292. Reprinted with permission.)

Appendix 5

Other useful disease prevention resources

- Guay DRP. Pet-assisted therapy in the nursing home setting: potential for zoonosis. *Am J Infect Control* 2001;29:178–186.
- Lefebvre SL, Golab GC, Christensen EL, et al. Guidelines for animal-assisted interventions in health care facilities. *Am J Infect Control* 2008;36:78–85.
- Murthy R, Bearman G, Brown S, et al. Animals in healthcare facilities: recommendations to minimize potential risks. *Infect Control Hosp Epidemiol* 2015;36:495–516.
- National Institute for Occupational Safety and Health. Veterinary Safety and Health. Available at: www.cdc.gov/niosh/topics/veterinary/default.html. Accessed Feb 9, 2017.
- Sehulster L, Chinn RYW. Guidelines for environmental infection control in health-care facilities: recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee (HIPAC) (Erratum published in *MMWR Recomm Rep* 2003;52:1025). *MMWR Recomm Rep* 2003;52:1–42.
- Varela K, Brown J, Lipton B, et al. A Review of Zoonotic Disease Threats to Pet Owners: A Compendium of Measures to Prevent Zoonotic Diseases Associated with Non-Traditional Pets Such as Rodents and Other Small Mammals, Reptiles, Amphibians, Backyard Poultry, and Other Selected Animals. *Vector-Borne and Zoonotic Diseases*, 2022; (22)303-360. <http://doi.org/10.1089/vbz.2022.0022>
- Williams CJ, Scheftel JM, Elchos BL, et al. Compendium of veterinary standard precautions for zoonotic disease prevention in veterinary personnel: National Association of State Public Health Veterinarians: Veterinary Infection Control Committee 2015 (Erratum published in *J Am Vet Med Assoc* 2016;248:171). *J Am Vet Med Assoc* 2015;247:1252– 1277.